

# The Binet formula, sums and representations of generalized Fibonacci $p$ -numbers

Emrah Kilic

*TOBB ETU University of Economics and Technology, Mathematics Department, 06560 Sogutozu, Ankara, Turkey*

Received 10 June 2006; accepted 6 March 2007

Available online 1 April 2007

---

## Abstract

In this paper, we consider the generalized Fibonacci  $p$ -numbers and then we give the generalized Binet formula, sums, combinatorial representations and generating function of the generalized Fibonacci  $p$ -numbers. Also, using matrix methods, we derive an explicit formula for the sums of the generalized Fibonacci  $p$ -numbers.

© 2007 Elsevier Ltd. All rights reserved.

---

## 1. Introduction

We consider a generalization of well-known Fibonacci numbers, which are called Fibonacci  $p$ -numbers. The Fibonacci  $p$ -numbers  $F_p(n)$  are defined by the following equation for  $n > p + 1$

$$F_p(n) = F_p(n - 1) + F_p(n - p - 1) \quad (1)$$

with initial conditions

$$F_p(1) = F_p(2) = \dots = F_p(p) = F_p(p + 1) = 1.$$

If we take  $p = 1$ , then the sequence of Fibonacci  $p$ -numbers,  $\{F_p(n)\}$ , is reduced to the well-known Fibonacci sequence  $\{F_n\}$ .

The Fibonacci  $p$ -numbers and their properties have been studied by some authors (for more details see [1,4–6,8,13–26,29]).

---

*E-mail address:* [ekilic@etu.edu.tr](mailto:ekilic@etu.edu.tr).

In 1843, Binet gave a formula which is called “Binet formula” for the usual Fibonacci numbers  $F_n$  by using the roots of the characteristic equation  $x^2 - x - 1 = 0 : \alpha = \frac{1+\sqrt{5}}{2}, \beta = \frac{1-\sqrt{5}}{2}$

$$F_n = \frac{\alpha^n - \beta^n}{\alpha - \beta}$$

where  $\alpha$  is called Golden Proportion,  $\alpha = \frac{1+\sqrt{5}}{2}$  (for details see [7,30,28]). In [12], Levesque gave a Binet formula for the Fibonacci sequence by using a generating function. In [2], the authors considered an  $n \times n$  companion matrix and its  $n$ th power, then gave the combinatorial representation of the sequence generated by the  $n$ th power the matrix. Further in [25], the authors derived analytical formulas for the Fibonacci  $p$ -numbers and then showed these formulas are similar to the Binet formulas for the classical Fibonacci numbers. Also, in [11], the authors gave the generalized Binet formulas and the combinatorial representations for the generalized order- $k$  Fibonacci [3] and Lucas [27] numbers. In [10], the authors defined the generalized order- $k$  Pell numbers and gave the Binet formula for the generalized Pell sequence. For the common generalization of the generalized order- $k$  Fibonacci and Pell numbers, and its generating matrix, sums and combinatorial representation, we refer readers to [9].

In this paper, we consider the generalized Fibonacci  $p$ -numbers and give the generalized Binet formula, combinatorial representations and sums of the generalized Fibonacci  $p$ -numbers by using the matrix method.

The generating matrix for the generalized Fibonacci  $p$ -numbers is given by Stakhov [23] as follows: Let  $Q_p$  be the following  $(p + 1) \times (p + 1)$  companion matrix :

$$Q_p = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 1 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ \vdots & \dots & \ddots & \dots & \vdots & \vdots \\ 0 & \dots & 0 & 1 & 0 & 0 \\ 0 & 0 & \dots & 0 & 1 & 0 \end{bmatrix} \tag{2}$$

and the  $n$ th power of the matrix  $Q_p$  is

$$Q_p^n = \begin{bmatrix} F_p(n+1) & F_p(n-p+1) & \dots & F_p(n-1) & F_p(n) \\ F_p(n) & F_p(n-p) & \dots & F_p(n-2) & F_p(n-1) \\ \vdots & \vdots & & \vdots & \vdots \\ F_p(n-p+2) & F_p(n-2p+2) & \dots & F_p(n-p) & F_p(n-p+1) \\ F_p(n-p+1) & F_p(n-2p+1) & \dots & F_p(n-p-1) & F_p(n-p) \end{bmatrix}. \tag{3}$$

The matrix  $Q_p$  is said to be a generalized Fibonacci  $p$ -matrix.

### 2. The generalized Binet formula

In this section, we give the generalized Binet formula for the generalized Fibonacci  $p$ -numbers. We start with the following results.

**Lemma 1.** Let  $a_p = \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1}$ . Then  $a_p > a_{p+1}$  for  $p > 1$ .

**Proof.** Since  $2p^3 - 2p - 1 > 0$  and  $p > 1$ ,  $(p^2 + 2p + 1)(p^2 - 1) > p^4$ . Thus,  $\left(\frac{p^2-1}{p^2}\right) > \left(\frac{p}{p+1}\right)^2$ . Therefore, for  $p > 1$ ,  $\left(\frac{p^2-1}{p^2}\right)^{p-1} > \left(\frac{p}{p+1}\right)^2$  and so  $\left(\left(\frac{p-1}{p^2}\right) \times \left(\frac{p+1}{p}\right)\right)^{p-1} > \left(\frac{p}{p+1}\right)^2$ . Then we have  $\left(\frac{p-1}{p^2}\right)^{p-1} > \left(\frac{p}{p+1}\right)^{p+1}$ . So the proof is easily seen.  $\square$

**Lemma 2.** *The characteristic equation of the Fibonacci  $p$ -numbers  $x^p - x^{p-1} - 1 = 0$  does not have multiple roots for  $p > 1$ .*

**Proof.** Let  $f(z) = z^p - z^{p-1} - 1$ . Suppose that  $\alpha$  is a multiple root of  $f(z) = 0$ . Note that  $\alpha \neq 0$  and  $\alpha \neq 1$ . Since  $\alpha$  is a multiple root,  $f(\alpha) = \alpha^p - \alpha^{p-1} - 1 = 0$  and  $f'(\alpha) = p\alpha^{p-1} - (p-1)\alpha^{p-2} = 0$ . Then

$$f'(\alpha) = \alpha^{p-2}(p\alpha - (p-1)) = 0.$$

Thus  $\alpha = \frac{p-1}{p}$ , and hence

$$\begin{aligned} 0 &= f(\alpha) = -\alpha^p + \alpha^{p-1} + 1 = \alpha^{p-1}(1 - \alpha) + 1 \\ &= \left(\frac{p-1}{p}\right)^{p-1} \left(1 - \frac{p-1}{p}\right) + 1 = \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1} + 1 \\ &= a_p + 1. \end{aligned}$$

Since, by Lemma 1,  $a_2 = \frac{1}{4} < 1$  and  $a_p > a_{p+1}$  for  $p > 1$ ,  $a_p \neq 1$ , which is a contradiction. Therefore, the equation  $f(z) = 0$  does not have multiple roots.  $\square$

We suppose that  $f(\lambda)$  is the characteristic polynomial of the generalized Fibonacci  $p$ -matrix  $Q_p$ . Then,  $f(\lambda) = \lambda^{p+1} - \lambda^p - 1$ , which is a well-known fact from the companion matrices. Let  $\lambda_1, \lambda_2, \dots, \lambda_{p+1}$  be the eigenvalues of the matrix  $Q_p$ . Then, by Lemma 2, we know that  $\lambda_1, \lambda_2, \dots, \lambda_{p+1}$  are distinct. Let  $A$  be a  $(p+1) \times (p+1)$  Vandermonde matrix as follows:

$$A = \begin{bmatrix} \lambda_1^p & \lambda_1^{p-1} & \dots & \lambda_1 & 1 \\ \lambda_2^p & \lambda_2^{p-1} & \dots & \lambda_2 & 1 \\ \vdots & \vdots & & \vdots & \vdots \\ \lambda_{p+1}^p & \lambda_{p+1}^{p-1} & \dots & \lambda_{p+1} & 1 \end{bmatrix}.$$

We denote  $A^T$  by  $V$ . Let

$$d_k^i = \begin{bmatrix} \lambda_1^{n+p+1-i} \\ \lambda_2^{n+p+1-i} \\ \vdots \\ \lambda_{p+1}^{n+p+1-i} \end{bmatrix}$$

and  $V_j^{(i)}$  be a  $(p+1) \times (p+1)$  matrix obtained from  $V$  by replacing the  $j$ th column of  $V$  by  $d_k^i$ .

Then we can give the generalized Binet formula for the generalized Fibonacci  $p$ -numbers with the following theorem.

**Theorem 3.** Let  $F_p(n)$  be the  $n$ th generalized Fibonacci  $p$ -number; then

$$q_{ij} = \frac{\det(V_j^{(i)})}{\det(V)}$$

where  $Q_p^n = [q_{ij}]$  and  $q_{ij} = F_p(n + j - i - p)$  for  $j \geq 2$  and  $q_{i,1} = F_p(n + 2 - i)$  for  $j = 1$ .

**Proof.** Since the eigenvalues of the matrix  $Q_p$  are distinct, the matrix  $Q_p$  is diagonalizable. It is easy to show that  $Q_p V = V D$ , where  $D = \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_{p+1})$ . Since the Vandermonde matrix  $V$  is invertible,  $V^{-1} Q_p V = D$ . Hence, the matrix  $Q_p$  is similar to the diagonal matrix  $D$ . So we have the matrix equation  $Q_p^n V = V D^n$ . Since  $Q_p^n = [q_{ij}]$ , we have the following linear system of equations:

$$\begin{aligned} q_{i1}\lambda_1^p + q_{i2}\lambda_1^{p-1} + \dots + q_{i,p+1} &= \lambda_1^{p+n+1-i} \\ q_{i1}\lambda_2^p + q_{i2}\lambda_2^{p-1} + \dots + q_{i,p+1} &= \lambda_2^{p+n+1-i} \\ &\vdots \\ q_{i1}\lambda_{p+1}^p + q_{i2}\lambda_{p+1}^{p-1} + \dots + q_{i,p+1} &= \lambda_{p+1}^{p+n+1-i}. \end{aligned}$$

Thus, for each  $j = 1, 2, \dots, p + 1$ , we obtain

$$q_{ij} = \frac{\det(V_j^{(i)})}{\det(V)}.$$

So the proof is complete.  $\square$

Thus, we give the Binet formula for the  $n$ th Fibonacci  $p$ -number  $F_p(n)$  by the following corollary.

**Corollary 4.** Let  $F_p(n)$  be the  $n$ th Fibonacci  $p$ -number. Then

$$F_p(n) = \frac{\det(V_1^{(2)})}{\det(V)} = \frac{\det(V_{p+1}^{(1)})}{\det(V)}.$$

**Proof.** The conclusion is immediate result of Theorem 3 by taking  $i = 2, j = 1$  or  $i = 1, j = p + 1$ .  $\square$

The following lemma can be obtained from [2].

**Lemma 5.** Let the matrix  $Q_p^n = [q_{ij}]$  be as in (3). Then

$$q_{ij} = \sum_{(m_1, \dots, m_{p+1})} \frac{m_j + m_{j+1} + \dots + m_{p+1}}{m_1 + m_2 + \dots + m_{p+1}} \times \binom{m_1 + m_2 + \dots + m_{p+1}}{m_1, m_2, \dots, m_{p+1}}$$

where the summation is over nonnegative integers satisfying  $m_1 + 2m_2 + \dots + (p + 1)m_{p+1} = n - i + j$ , and defined to be 1 if  $n = i - j$ .

Then we have the following corollaries.

**Corollary 6.** Let  $F_p(n)$  be the generalized Fibonacci  $p$ -number. Then

$$F_p(n) = \sum_{(m_1, \dots, m_{p+1})} \frac{m_{p+1}}{m_1 + m_2 + \dots + m_{p+1}} \times \binom{m_1 + m_2 + \dots + m_{p+1}}{m_1, m_2, \dots, m_{p+1}}$$

where the summation is over nonnegative integers satisfying  $m_1 + 2m_2 + \dots + (p + 1)m_{p+1} = n + p$ .

**Proof.** In Lemma 5, when  $i = 1$  and  $j = p + 1$ , then the conclusion can be directly seen from (3).  $\square$

**Corollary 7.** Let  $F_p(n)$  be the generalized Fibonacci  $p$ -number. Then

$$F_p(n) = \sum_{(m_1, \dots, m_{p+1})} \binom{m_1 + m_2 + \dots + m_{p+1}}{m_1, m_2, \dots, m_{p+1}}$$

where the summation is over nonnegative integers satisfying  $m_1 + 2m_2 + \dots + (p + 1)m_{p+1} = n - 1$ .

**Proof.** In Lemma 5, if we take  $i = 2$  and  $j = 1$ , then we have the corollary from (3).  $\square$

We consider the generating function of the generalized Fibonacci  $p$ -numbers. We give the following lemma.

**Lemma 8.** Let  $F_p(n)$  be the  $n$ th generalized Fibonacci number, then for  $n > 1$

$$x^n = F_p(n - p + 1)x^p + \sum_{j=1}^p F_p(n - p + 1 - j) x^{j-1}.$$

**Proof.** We suppose that  $n = p + 1$ ; then by the definition of the Fibonacci  $p$ -numbers

$$x^{p+1} = F_p(2)x^p + F_p(1) = x^p + 1.$$

Now we suppose that the equation holds for any integer  $n, n > p + 1$ . Then we show that the equation holds for  $n + 1$ . Thus, from our assumption and the characteristic equation the Fibonacci  $p$ -numbers,

$$\begin{aligned} x^{n+1} &= x^n x = \left( F_p(n - p + 1)x^p + \sum_{j=1}^p F_p(n - p + 1 - j) x^{j-1} \right) x \\ &= F_p(n - p + 1) (x^p + 1) + \sum_{j=1}^p F_p(n - p + 1 - j) x^j \\ &= F_p(n - p + 1)x^p + F_p(n - p + 1) + F_p(n - 2p + 1) x^p \\ &\quad + F_p(n - 2p + 2)x^{p-1} + \dots + F_p(n - 2p + 1)x^2 + F_p(n - p)x \\ &= [F_p(n - p + 1) + F_p(n - 2p + 1)] x^p + F_p(n - 2p + 2)x^{p-1} \\ &\quad + F_p(n - 2p + 3) x^{p-2} + \dots + F_p(n - p)x + F_p(n - p + 1). \end{aligned} \tag{4}$$

Using the definition of the generalized Fibonacci  $p$ -numbers, we have

$$F_p(n - p + 1) + F_p(n - 2p + 1) = F_p(n - p + 2).$$

Therefore, we can write the Eq. (4) as follows

$$\begin{aligned} x^{n+1} &= F_p(n - p + 2)x^p + F_p(n - 2p + 2)x^{p-1} \\ &\quad + F_p(n - 2p + 3)x^{p-2} + \dots + F_p(n - p)x + F_p(n - p + 1) \\ &= F_p(n - p + 2)x^p + \sum_{j=1}^p F_p(n - p + 2 - j)x^{j-1} \end{aligned} \tag{5}$$

which is what was desired.  $\square$

Now we give the generating function of the generalized Fibonacci  $p$ -numbers:

Let

$$G_p(x) = F_p(1) + F_p(2)x + F_p(3)x^2 + \dots + F_p(n + 1)x^n + \dots .$$

Then

$$G_p(x) - xG_p(x) - x^{p+1}G_p(x) = (1 - x - x^{p+1})G_p(x).$$

By the Eq. (5), we have  $(1 - x - x^{p+1})G_p(x) = F_p(1) = 1$ . Thus

$$G_p(x) = (1 - x - x^{p+1})^{-1}$$

for  $0 \leq x + x^{p+1} < 1$ .

Let  $f_p(x) = x + x^{p+1}$ . Then, for  $0 \leq f_p(x) < 1$ , we have the following lemma.

**Lemma 9.** For positive integers  $t$  and  $n$ , the coefficient of  $x^n$  in  $(f_p(x))^t$  is

$$\sum_{j=0}^t \binom{t}{j}, \quad \frac{n}{p+1} \leq t \leq n$$

where the integers  $j$  satisfy  $pj + t = n$ .

**Proof.** From the above results, we write

$$(f_p(x))^t = (x + x^{p+1})^t = x^t (1 + x^p)^t = x^t \sum_{j=0}^t \binom{t}{j} x^{pj}.$$

In the above equation, we consider the coefficient of  $x^n$ . For positive integers  $t$  and  $j$  such that  $pj + t = n$  and  $j \leq t$ , the coefficients of  $x^n$  are

$$\sum_{j=0}^t \binom{t}{j}, \quad \frac{n}{p+1} \leq t \leq n.$$

So we have the required conclusion.  $\square$

Now we can give a representation for the generalized Fibonacci  $p$ -numbers by the following theorem.

**Theorem 10.** Let  $F_p(n)$  be the  $n$ th generalized Fibonacci  $p$ -number. Then, for positive integers  $t$  and  $n$ ,

$$F_p(n+1) = \sum_{\frac{n}{p+1} \leq t \leq n} \sum_{j=0}^t \binom{t}{j}$$

where the integers  $j$  satisfy  $pj + t = n$ .

**Proof.** Since

$$\begin{aligned} G_p(x) &= F_p(1) + F_p(2)x + F_p(3)x^2 + \cdots + F_p(n+1)x^n + \cdots \\ &= \frac{1}{1-x-x^{p+1}} \end{aligned}$$

and  $f_p(x) = x + x^{p+1}$ , the coefficient of  $x^n$  is the  $(n+1)$ th generalized Fibonacci  $p$ -number,  $F_p(n+1)$  in  $G_p(x)$ . Thus

$$\begin{aligned} G_p(x) &= \frac{1}{1-x-x^{p+1}} \\ &= \frac{1}{1-f_p(x)} \\ &= 1 + f_p(x) + (f_p(x))^2 + \cdots + (f_p(x))^n + \cdots \\ &= 1 + x(1+x^p) + x^2 \sum_{j=0}^2 \binom{2}{j} x^{pj} + \cdots + x^n \sum_{j=0}^n \binom{n}{j} x^{pj} + \cdots \end{aligned}$$

As we need the coefficient of  $x^n$ , we only consider the first  $n+1$  terms on the right-side. Thus by [Lemma 9](#), the proof is complete.  $\square$

Now we give an exponential representation for the generalized Fibonacci  $p$ -numbers.

$$\begin{aligned} \ln G_p(x) &= \ln \left[ 1 - (x + x^{p+1}) \right]^{-1} \\ &= -\ln \left[ 1 - (x + x^{p+1}) \right] \\ &= - \left[ - (x + x^{p+1}) - \frac{1}{2} (x + x^{p+1})^2 - \cdots - \frac{1}{n} (x + x^{p+1})^n - \cdots \right] \\ &= x \left[ (1 + x^p) + \frac{1}{2} (1 + x^p)^2 + \cdots + \frac{1}{n} (1 + x^p)^n + \cdots \right] \\ &= x \sum_{n=0}^{\infty} \frac{1}{n} (1 + x^p)^n. \end{aligned}$$

Thus,

$$G_p(x) = \exp \left( x \sum_{n=0}^{\infty} \frac{1}{n} (1 + x^p)^n \right).$$

### 3. Sums of the generalized Fibonacci $p$ -numbers by matrix methods

In this section, we define a  $(p + 2) \times (p + 2)$  matrix  $T$ , and then we show that the sums of the generalized Fibonacci  $p$ -numbers can be obtained from the  $n$ th power of the matrix  $T$ .

**Definition 11.** For  $p \geq 1$ , let  $T = (t_{ij})$  denote the  $(p + 2) \times (p + 2)$  matrix by  $t_{11} = t_{21} = t_{22} = t_{2,p+2} = 1, t_{i+1,i} = 1$  for  $2 \leq i \leq p + 1$  and 0 otherwise.

Clearly, by the definition of the matrix  $Q_p$ ,

$$T = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & & \ddots & \vdots & \vdots \\ 0 & 0 & & 0 & 1 & 0 \end{bmatrix} \quad \text{or} \quad T = \begin{bmatrix} 1 & 0 & \dots & 0 \\ 1 & & & \\ 0 & Q_p & & \\ \vdots & & & \\ 0 & & & \end{bmatrix} \tag{6}$$

where the  $(p + 1) \times (p + 1)$  matrix  $Q_p$  given by (2).

Let  $S_n$  denote the sums of the generalized Fibonacci  $p$ -numbers from 1 to  $n$ , that is:

$$S_n = \sum_{i=1}^n F_p(i). \tag{7}$$

Now we define a  $(p + 2) \times (p + 2)$  matrix  $C_n$  as follows

$$C_n = \begin{bmatrix} 1 & 0 & \dots & 0 \\ S_n & & & \\ S_{n-1} & Q_p^n & & \\ \vdots & & & \\ S_{n-p} & & & \end{bmatrix} \tag{8}$$

where  $Q_p^n$  given by (3).

Then we have the following theorem.

**Theorem 12.** Let the  $(p + 2) \times (p + 2)$  matrices  $T$  and  $C_n$  be as in (6) and (8), respectively. Then, for  $n \geq 1$ :

$$C_n = T^n.$$

**Proof.** We will use the induction method to prove that  $C_n = T^n$ . If  $n = 1$ , then, by the definition of the matrix  $C_n$  and generalized Fibonacci  $p$ -numbers, we have

$$C_1 = T.$$

Now we suppose that the equation holds for  $n$ . Then we show that the equation holds for  $n + 1$ . Thus,

$$T^{n+1} = T^n \cdot T$$

and by our assumption,

$$T^{n+1} = C_n T.$$



Since  $S_{n+1} = S_n + F_p(n + 1)$  and using the definition of the generalized Fibonacci numbers, we can derive the following matrix recurrence relation

$$C_n T = C_{n+1}.$$

So the proof is complete.  $\square$

We define two  $(p + 2) \times (p + 2)$  matrices. First, we define the matrix  $R$  as follows:

$$R = \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ -1 & \lambda_1^p & \lambda_2^p & \dots & \lambda_{p+1}^p \\ -1 & \lambda_1^{p-1} & \lambda_2^{p-1} & \dots & \lambda_{p+1}^{p-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -1 & \lambda_1 & \lambda_2 & \dots & \lambda_{p+1} \\ -1 & 1 & 1 & \dots & 1 \end{bmatrix} \tag{9}$$

and the diagonal matrix  $D_1$  as follows:

$$D_1 = \begin{bmatrix} 1 & & & & \\ & \lambda_1 & & & \\ & & \ddots & & \\ & & & \ddots & \\ & & & & \lambda_{p+1} \end{bmatrix} \tag{10}$$

where the  $\lambda_i$ 's are the eigenvalues of the matrix  $Q_p$  for  $1 \leq i \leq p + 1$ .

We give the following theorem for the computing the sums of the generalized Fibonacci  $p$ -numbers 1 from to  $n$  by using a matrix method.

**Theorem 13.** *Let the sums of the generalized Fibonacci numbers  $S_n$  be as in (7). Then*

$$S_n = F_p(n + p + 1) - 1.$$

**Proof.** If we compute the  $\det R$  by the Laplace expansion of determinant with respect to the first row, then we obtain that  $\det R = \det V$ , where the Vandermonde matrix  $V$  is as in Theorem 3. Therefore, we can easily find the eigenvalues of the matrix  $R$ . Since the characteristic equation of the matrix  $R$  is  $(x^p - x^{p-1} - 1) \times (x - 1)$  and by Lemma 2, the eigenvalues of the matrix  $R$  are  $1, \lambda_1, \dots, \lambda_{p+1}$  and distinct. So the matrix  $R$  is diagonalizable. We can easily prove that  $TR = RD_1$ , where the matrices  $T, R$  and  $D_1$  are as in (6), (9) and (10), respectively. Then we have

$$T^n R = R D_1^n. \tag{11}$$

Since  $T^n = C_n$ , we write that  $C_n R = R D_1^n$ . We know that  $S_n = (C_n)_{2,1}$ . By a matrix multiplication,

$$S_n - \left( \sum_{i=0}^p F_p(n + 1 - i) \right) = -1. \tag{12}$$

By the definition of the generalized Fibonacci  $p$ -numbers, we know that  $\sum_{i=0}^p F_p(n + 1 - i) = F_p(n + p + 1)$ . Then we write the Eq. (12) as follows:

$$S_n - F_p(n + p + 1) = -1.$$




Fig. 1.

Thus,

$$S_n = \sum_{i=1}^n F_p(i) = F_p(n+p+1) - 1.$$

So the proof is complete.  $\square$

In [30], the author presents an enumeration problem for the paths from  $A$  to  $c_n$ , and then shows that the number of paths from  $A$  to  $c_n$  are equal to the  $n$ th usual Fibonacci number. Now, we are interested in a problem of paths. The problem is as in Fig. 1.

It is seen that the number of path from  $A$  to  $c_1, c_2, \dots, c_{p+1}$  is 1. Also, we know that the initial conditions of the generalized Fibonacci  $p$ -numbers, that is,  $F_p(1), F_p(2), \dots, F_p(p+1)$ , are 1. Now we consider the case  $n > p+1$ . The number of the path from  $A$  to  $c_{p+2}$  is 2. By the induction method, one can see that the number of the path from  $A$  to  $c_n$  is the  $n$ th generalized Fibonacci  $p$ -number.

## References

- [1] B.A. Bondarenko, Generalized Pascal's Triangles and Pyramids: Their fractals, Graphs, and Applications, Fibonacci Association, 1993.
- [2] W.Y.C. Chen, J.D. Louck, The combinatorial power of the companion matrix, *Linear Algebra Appl.* 232 (1996) 261–278.
- [3] M.C. Er, Sums of Fibonacci numbers by matrix methods, *Fibonacci Quart.* 22 (3) (1984) 204–207.
- [4] S. Falcon, A. Plaza, The  $k$ -Fibonacci hyperbolic functions, *Chaos Solitons Fractals*, doi:10.1016/j.chaos.2006.11.019.
- [5] S. Falcon, A. Plaza, The  $k$ -Fibonacci sequence and the Pascal 2-Triangle, *Chaos Solitons Fractals* 33 (1) (2007) 38–49.
- [6] M.J.G. Gazale, From Pharaons to Fractals, Princeton University Press, Princeton, New Jersey, 1999 (Russian translation, 2002).
- [7] V.E. Hoggat, Fibonacci and Lucas Numbers, Houghton-Mifflin, PaloAlto, California, 1969.
- [8] J. Kappraff, Connections. The Geometric Bridge Between Art and Science, second ed., World Scientific, Singapore, New Jersey, London, Hong Kong, 2001.
- [9] E. Kilic, The generalized order- $k$  Fibonacci-Pell sequence by matrix methods, doi:10.1016/j.cam.2006.10.071.
- [10] E. Kilic, D. Tasci, The generalized Binet formula, representation and sums of the generalized order- $k$  Pell numbers, *Taiwanese J. Math.* 10 (6) (2006) 1661–1670.
- [11] E. Kilic, D. Tasci, On the generalized order- $k$  Fibonacci and Lucas numbers, *Rocky Mountain J. Math.* 36 (6) (2006) 1915–1926.
- [12] C. Levesque, On  $m$ th-order linear recurrences, *Fibonacci Quart.* 23 (4) (1985) 290–293.
- [13] B. Rozin, The Golden Section: A morphological law of living matter, Available from [www.goldensection.net](http://www.goldensection.net).
- [14] N.A. Soljanichenko, B.N. Rozin, Mystery of Golden Section, Theses of the Conference Fenid-90: Non-traditional ideas about Nature and its phenomena, (3) Homel, 1990.
- [15] C.P. Spears, M. Bicknell-Johnson, Asymmetric Cell Dision: Binomial Identities for Age Analysis of Mortal vs. Immortal Trees, in: Applications of Fibonacci Numbers, vol. 7, 1998, pp. 377–391.

- [16] A.P. Stakhov, Introduction into Algorithmic Measurement Theory, Soviet Radio, Moscow, 1977.
- [17] A.P. Stakhov, B. Rozin, The golden shofar, *Chaos Solitions Fractals* 26 (3) (2005) 677–684.
- [18] A.P. Stakhov, Algorithmic measurement theory, in: *Mathematics & Cybernetics*, vol. 6, Moscow, 1979.
- [19] A.P. Stakhov, Codes of the golden proportion, *Radio Commun.* (1984).
- [20] A.P. Stakhov, The golden section in the measurement theory, *Comput. Math. Appl.* 17 (4–6) (1989) 613–638.
- [21] A.P. Stakhov, I.S. Tkachenko, Hyperbolic fibonacci trigonometry, *Rep. Ukrainian Acad. Sci.* 208 (7) (1993) 9–14.
- [22] A.P. Stakhov, The Golden Section and Modern Harmony Mathematics, in: *Applications of Fibonacci numbers*, vol. 7, Kluwer Academic Publisher, 1998, pp. 393–399.
- [23] A.P. Stakhov, A generalization of the Fibonacci  $Q$ -matrix, *Rep. Natl. Acad. Sci. Ukraine* 9 (1999) 46–49.
- [24] A.P. Stakhov, B. Rozin, The continuous functions for the Fibonacci and Lucas  $p$ -numbers, *Chaos Solitions Fractals* 28 (2006) 1014–1025.
- [25] A.P. Stakhov, B. Rozin, Theory of Binet formulas for Fibonacci and Lucas  $p$ -numbers, *Chaos Solitions Fractals* 27 (2006) 1162–1177.
- [26] A.P. Stakhov, A.A. Sluchenkova, V. Massingua, Introduction into Fibonacci Coding and Cryptography, Osnova, Kharkov, 1999.
- [27] D. Tasci, E. Kilic, On the order- $k$  generalized Lucas numbers, *Appl. Math. Comput.* 155 (3) (2004) 637–641.
- [28] S. Vajda, Fibonacci and Lucas numbers, and the Golden Section, Theory and applications, John Wiley & Sons, New York, 1989.
- [29] V.W. de Spinadel, From the golden mean to chaos, in: *Nueva Libreria*, 1998. second edition Nobuko, 2004.
- [30] N.N. Vorob'ev, Fibonacci Numbers, Nauka, Moscow, 1978.