



On the Fibonacci and Lucas p -numbers, their sums, families of bipartite graphs and permanents of certain matrices

E. Kilic *, A.P. Stakhov

*TOBB Economics and Technology University, Mathematics Department, Sogutozu , 06560 Ankara, Turkey
International Club of the Golden Section, 6 McCreary Trail, Bolton, Ont., Canada L7E 2C8*

Accepted 4 October 2007

Abstract

In this paper we consider certain generalizations of the well-known Fibonacci and Lucas numbers, the generalized Fibonacci and Lucas p -numbers. We give relationships between the generalized Fibonacci p -numbers, $F_p(n)$, and their sums, $\sum_{i=1}^n F_p(i)$, and the 1-factors of a class of bipartite graphs. Further we determine certain matrices whose permanents generate the Lucas p -numbers and their sums.

© 2007 Elsevier Ltd. All rights reserved.

1. Introduction

The well-known Fibonacci $\{F_n\}$ and Lucas $\{L_n\}$ sequences are defined by the following equations, for $n > 1$

$$F_n = F_{n-1} + F_{n-2} \quad (1)$$

where $F_0 = 0$, $F_1 = 1$, and

$$L_n = L_{n-1} + L_{n-2} \quad (2)$$

where $L_0 = 2$, $L_1 = 1$.

Modern science, particularly physics, [23,24,28–33,57], widely applies these sequences $\{F_n\}$ and $\{L_n\}$ which result from application of the following recurrence relations: for $n > 1$

$$F(n) = F(n-1) + F(n-2) \quad (3)$$

$$F(0) = 0, \quad F(1) = 1$$

$$L(n) = L(n-1) + L(n-2) \quad (4)$$

$$L(0) = 2, \quad L(1) = 1$$

In [46,48], the authors considered the rules (3) and (4), then gave the generalization of the Fibonacci and Lucas numbers, called the Fibonacci and Lucas p -numbers as, for any given $p(p = 1, 2, 3, \dots)$ and $n > p + 1$

* Corresponding author.

E-mail address: ekilic@etu.edu.tr (E. Kilic).

$$F_p(n) = F_p(n - 1) + F_p(n - p - 1) \tag{5}$$

with initial conditions $F_p(0) = 0, F_p(1) = \dots = F_p(p) = F_p(p + 1) = 1$, and for $n > p$

$$L_p(n) = L_p(n - 1) + L_p(n - p - 1) \tag{6}$$

where $L_p(0) = p + 1, L_p(1) = \dots = L_p(p) = 1$, respectively.

The ratio of the successive Fibonacci number is as known *golden ratio*, that is

$$\tau = \frac{F_{n+1}}{F_n}$$

which is approximately equal 1.618. Also the golden ratio satisfies the characteristic equation of the Fibonacci numbers $t^2 - t - 1 = 0$. When the case of the Fibonacci p -numbers, the well-known golden ratio is called p -*proportion* and shown by τ_p which satisfies the following equation:

$$t^{p+1} = t^p + 1$$

There are many applications of the golden ratio and p -proportion in many places of mathematics and physics. For example, we can take the values of the resistors in Fig. 1 as follows:

$$R1 = \tau_p^{-p}R; \quad R2 = \tau_p^{p+1}R; \quad R3 = \tau_p R$$

where τ_p is the golden p -proportion, $p \in \{0, 1, 2, 3, \dots\}$.

It is clear that the divider in Fig. 1 gives an infinite number of the different resistor dividers because every p originates a new divider. In particular, for the case $p = 0$ the value of the golden 0-proportion $\tau_0 = 2$ and the divider is reduced to the classical *binary* divider. For the case $p = 1$ the resistors $R1, R2, R3$ take the following values:

$$R1 = \tau^{-1}R; \quad R2 = \tau^2R; \quad R3 = \tau R$$

where $\tau = (1 + \sqrt{5})/2$ is the classical golden mean. More details can be found in [40].

Furthermore, as examples of physical applications of the golden ratio, in a general theory of high energy particle theory, i.e., the golden mean have been widely used. For other applications of the golden mean, we can refer to the well-known works of El Naschie and Marek-Crnjac [20–22,34–39].

Recently, in [40–53,58], many interesting properties and applications of these recurrences have been studied by several authors. Especially, in [50,52,53], Stakhov gave the generating matrices of the Fibonacci p -numbers called “golden” matrices and their inverses. Then the author gave the interesting applications to the coding theory called “golden” cryptography. Also in [47], one can find the Binet type formulas for these recurrences and many interesting properties. In [16], considering the generating matrix of the sequence of the Fibonacci p -numbers, the author used the matrix methods and then gave the Binet formula, sums and combinatorial representations of the Fibonacci p -numbers. One can find many properties of the Fibonacci and Lucas p -numbers in webpage of the Museum of Harmony and Golden Section “<http://www.goldenmuseum.com>”.

Further in the earlier works, one can find another generalizations of the Fibonacci and Lucas numbers. For example, in [25], the author defined the k -generalized Fibonacci numbers as

$$f_n = \sum_{j=1}^k f_{n-j} \quad \text{for } n > k \geq 2 \tag{7}$$

with

$$f_0 = f_1 = \dots = f_{k-2} = 0, \quad f_{k-1} = f_k = 1$$

where f_n is the n th k -generalized Fibonacci number.




Fig. 1.

In [4], Er considered the definition of k -generalized Fibonacci number, and then defined k sequences of generalized order- k Fibonacci numbers as shown:

$$g_n^i = \sum_{j=1}^k g_{n-j}^i \quad \text{for } n > 0 \quad \text{and } 1 \leq i \leq k \tag{8}$$

with initial conditions

$$g_n^i = \begin{cases} 1 & \text{if } n = 1 - i, \\ 0 & \text{otherwise} \end{cases} \quad 1 - k \leq n \leq 0 \tag{9}$$

where g_n^i is the n th term of the i th sequence.

In [56], the authors defined k sequences of generalized order- k Lucas numbers as shown:

$$l_n^i = \sum_{j=1}^k l_{n-j}^i \quad \text{for } n > 0 \quad \text{and } 1 \leq i \leq k \tag{10}$$

with initial conditions

$$l_n^i = \begin{cases} -1 & \text{if } n = 1 - i, \\ 2 & \text{if } n = 2 - i, 1 - k \leq n \leq 0 \\ 0 & \text{otherwise} \end{cases}$$

where l_n^i is the n th term of the i th sequence. For the relationships between the generalized order- k Fibonacci, Lucas numbers and their Binet formulas and combinatorial representations see [10].

Let A and B be nonzero, relatively prime integers such that $D = A^2 - 4B \neq 0$. Define the generalized Fibonacci sequence, $\{u_n\}$, and the generalized Lucas sequence, $\{v_n\}$, by for all $n \geq 2$

$$u_n = Au_{n-1} - Bu_{n-2} \tag{11}$$

$$v_n = Av_{n-1} - Bv_{n-2} \tag{12}$$

where $u_0 = 0, u_1 = 1$ and $v_0 = 2, v_1 = A$. If $A = 1$ and $B = -1$, then $u_n = F_n$ (the n th Fibonacci number) and $v_n = L_n$ (the n th Lucas number).

As a special case of the sequence $\{u_n\}$, in [5], the authors consider k -Fibonacci numbers and their some properties. The *permanent* of an n -square matrix $A = (a_{ij})$ is defined by

$$\text{per } A = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i\sigma(i)}$$

where the summation extends over all permutations σ of the symmetric group S_n . The most important applications of permanents are in the areas of physics and chemistry. One can find more applications of permanents in [27].

The permanent of a matrix is analogous to the determinant, where all of the signs used in the Laplace expansion of minors are positive.

A matrix is said to be a $(0, 1)$ -matrix if each of its entries 0 or 1.

Many connections between permanents or determinants of tridiagonal matrices and the Fibonacci, Lucas numbers can be found in literature. For example, Minc [26] defined an $n \times n$ super-diagonal $(0, 1)$ -matrix $F(n, k)$ for $n + 1 \geq k$, and showed that the permanent of $F(n, k)$ equals a generalized order- k Fibonacci number. When $k = 2$, the matrix $F(n, 2)$ is reduced to the tridiagonal matrix and its permanent equals a usual Fibonacci number. Also in [54,55], the authors defined a family of tridiagonal matrices $M(n)$ and showed that the determinants of $M(n)$ are the Fibonacci numbers F_{2n+2} . In [19], Lehmer discussed the relationships between permanents of tridiagonal matrices, recurrence relations, and continued fractions. In [9], the authors defined two tridiagonal matrices and then gave the relationships of the permanents and determinants of these matrices and the second order linear recurrences given by (11) and (12). In [11], the authors present a result involving the permanent of an $(-1, 0, 1)$ -matrix and the Fibonacci number F_{n+1} . The authors then explore similar directions involving the positive subscripted Fibonacci and Lucas Numbers as well as their uncommon negatively subscripted counterparts. For further similar relationships, we can refer to [2,3,12,13,15,17,18].

Let $A = [a_{ij}]$ be an $m \times n$ real matrix having row vectors $\alpha_1, \alpha_2, \dots, \alpha_m$. We say that A is *contractible on column* (resp. *row*) k if *column* (resp. *row*) k contains exactly two nonzero entries. Suppose A is contractible on column k with $a_{ik} \neq 0 \neq a_{jk}$ and $i \neq j$. Then the $(m - 1) \times (n - 1)$ matrix $A_{ij:k}$ obtained from A by replacing row i with $a_{jk}\alpha_i + a_{ik}\alpha_j$ and deleting row j and column k is called *the contraction of A on column k relative to rows i and j* . If A is contractible on row k with $a_{ki} \neq 0 \neq a_{kj}$ and $i \neq j$, then the matrix $A_{k:ij} = [A_{ij:k}]^T$ is called *the contraction of A on row k relative to*

$$M^{(1)}(n,p) = \begin{pmatrix} 1 & 0 & \dots & 0 & 1 & 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & 1 & 0 & \dots \\ 0 & 1 & 1 & 0 & \dots & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & \dots & \dots & 0 & 1 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & \dots & 0 & 1 & 1 & 0 & \dots \\ 0 & 0 & 0 & \dots & \dots & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & \dots & 0 & 1 & 1 \end{pmatrix}$$

Since the matrix $M^{(1)}(n,p)$ can be contracted on column 1 and $F_p(0) = 0, F_p(1) = F_p(2) = F_p(3) = 1,$

$$M^{(2)}(n,p) = \begin{pmatrix} 1 & 0 & \dots & 0 & 1 & 1 & 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 0 & 0 & 1 & 0 & \dots \\ 0 & 1 & 1 & 0 & \dots & 0 & 0 & 0 & 1 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & \dots & \dots & \dots & \dots & 0 & 0 & 1 & 1 \end{pmatrix}$$

where $m_{1,p-1}^{(2)} = F_p(3), m_{1,p}^{(2)} = F_p(2), m_{1,p+1}^{(2)} = F_p(1).$ Continuing this process and since $F_p(p+1) = F_p(p) = \dots = F_p(1) = 1, F_p(0) = 0,$ we obtain

$$M^{(p-1)}(n,p) = \begin{pmatrix} 1 & 1 & \dots & 1 & 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \dots \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & 0 & 1 & 1 & 0 & 0 \\ 0 & \dots & 0 & \dots & 0 & 1 & 1 & 0 \\ 0 & \dots & \dots & 0 & \dots & 0 & 1 & 1 \end{pmatrix}$$

where $m_{1,1}^{(p-1)} = F_p(p+1), m_{1,2}^{(p-1)} = F_p(p), \dots, m_{1,p}^{(p-1)} = F_p(2), m_{1,p+1}^{(p-1)} = F_p(p).$

Now we consider the case $p \leq k \leq n - 4.$ Since the matrix $M^{(p-1)}(n,p)$ can be contracted on column 1 and $F_p(p+2) = 2$

$$M^{(p)}(n,p) = \begin{pmatrix} 2 & 1 & \dots & 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 \end{pmatrix}$$

where $m_{1,1}^{(p)} = F_p(p+2), m_{1,2}^{(p)} = F_p(p+1), \dots, m_{1,p}^{(p)} = F_p(3), m_{1,p+1}^{(p)} = F_p(p+1).$ Since the matrix $M^{(p)}(n,p)$ can be contracted on column 1

$$M^{(p+1)}(n,p) = \begin{pmatrix} 3 & 1 & \dots & 1 & 2 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \dots \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 0 & 1 & 1 & 0 & \dots \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{pmatrix}$$

where $m_{1,1}^{(p+1)} = F_p(p+3), m_{1,2}^{(p+1)} = F_p(p+2), \dots, m_{1,p}^{(p+1)} = F_p(4), m_{1,p+1}^{(p+1)} = F_p(p+2).$

We only consider the first row because the entries of other rows of $M^{(k)}(n,p)$ are 0 or 1 for all $k > 1$. Continuing this process, that is, by repeated contractions, we have

$$\begin{aligned} m_{1,1}^{(n-p-1)} &= F_p(n-p+1), \\ m_{1,2}^{(n-p-1)} &= F_p(n-2p+1), \\ m_{1,3}^{(n-p-1)} &= F_p(n-2p+2), \\ &\dots \\ m_{1,p+1}^{(n-p-1)} &= F_p(n-p). \end{aligned}$$

That is, the matrix $M^{(n-p-1)}(n,p)$ is as follows:

$$\begin{bmatrix} F_p(n-p+1) & F_p(n-2p+1) & \dots & \dots & \dots & F_p(n-p-1) & F_p(n-p) \\ 1 & 1 & 0 & \dots & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & \dots & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 1 & 1 \end{bmatrix}.$$

By repeating contractions

$$\begin{aligned} m_{11}^{(n-p)} &= F_p(n-p+1) + F_p(n-2p+1) = F_p(n-p+2) \\ m_{12}^{(n-p)} &= F_p(n-2p+2), \\ m_{13}^{(n-p)} &= F_p(n-2p+3) \\ &\vdots \\ m_{1,p}^{(n-p)} &= F_p(n-p) \end{aligned}$$

Thus, we have

$$M^{(n-3)}(n,p) = \begin{bmatrix} F_p(n-1) & F_p(n-p-1) & F_p(n-p-1) \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$$

Then we have, by contraction of $M^{(n-3)}(n,p)$ on column 1

$$M^{(n-2)}(n,p) = \begin{bmatrix} F_p(n) & F_p(n-p) \\ 1 & 1 \end{bmatrix}$$

By the Eq. (13), per $M^{(n-2)}(n,p) = \text{per } M(n,p) = F_p(n+1)$.

So the proof is complete. \square

For example, if we take $p = 2$, then we have, by [Theorem 2](#)

$$\text{per } M(n,2) = \text{per} \begin{bmatrix} 1 & 0 & 1 & 0 & \dots & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{bmatrix}_{n \times n} = F_2(n+1)$$

3. Sums of the consecutive generalized Fibonacci p -numbers

In this section, we determine a class of bipartite graph whose number of 1-factors is the sums of the consecutive generalized Fibonacci p -number, $\sum_{i=1}^n F_p(i)$.

Let n and p be positive integers such that $n > p \geq 1$.

Definition 3. Let $T(n, p) = [t_{ij}]$ be the $n \times n$ $(0, 1)$ -matrix with $t_{1,j} = 1$ for all j , $t_{i+1,i} = 1$ for $1 \leq i \leq n - 1$, $t_{i,i} = t_{i,i+p} = 1$ for $2 \leq i \leq n$ and a fixed integer p , and, 0 otherwise.

Clearly

$$T(n, p) = \begin{pmatrix} 1 & 1 & \dots & 1 & 1 & 1 & \dots & 1 & 1 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 & \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & \ddots & \ddots & \dots & \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{pmatrix}$$

Then we have the following Theorem.

Theorem 4. Let $G(T(n, p))$ be the bipartite graph with bipartite adjacency matrix $T(n, p)$, $n \geq 3$. Then the number of 1-factors of $G(T(n, p))$ is the sums of the consecutive generalized Fibonacci p -number, $\sum_{i=1}^n F_p(i)$.

Proof. We will use the induction method to prove that $per T(n, p) = \sum_{i=1}^n F_p(i)$. If $n = 3$, then we have, for $p = 2$

$$per T(3, 2) = per \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix} = \sum_{i=1}^3 F_2(i) = 3$$

If $n = 4$, then we have, for fixed $p = 2$

$$per T(4, 2) = per \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 0 & 1 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 1 & 0 \end{bmatrix} = \sum_{i=1}^4 F_2(i) = 5$$

Now we suppose that the equation holds for $n, n > p \geq 1$. If we compute the $per T(n, p)$ by the Laplace expansion of permanent with respect to the first column, then we have that

$$per T(n, p) = per \begin{bmatrix} 1 & 1 & \dots & 1 & 1 & 1 & \dots & 1 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \dots \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & \ddots & \ddots & \dots \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{bmatrix} + per \begin{bmatrix} 1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \\ 0 & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 & 0 & \dots & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & \ddots & \ddots & \dots \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{bmatrix}$$

Considering the definitions of the matrices $T(n, p)$ and $M(n, p)$, we write that

$$per T(n, p) = per T(n - 1, p) + per M(n - 1, p)$$

By our assumption and the [Theorem 2](#), we obtain that

$$\text{per } T(n, p) = \sum_{i=1}^n F_p(i) + F_p(n + 1) = \sum_{i=1}^{n+1} F_p(i)$$

So the proof is complete. \square

For example, when $p = 1$, then the sequence $\{F_p(n)\}$ is reduced to the well-known usual Fibonacci sequence $\{F_n\}$, then by [Theorem 4](#)

$$\text{per} \begin{bmatrix} 1 & 1 & 1 & \dots & 1 & 1 & 1 \\ 1 & 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & \dots & 0 \\ \vdots & 0 & 1 & 1 & 1 & \ddots & \vdots \\ \dots & \dots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & \dots & 0 & 1 & 1 & 1 \\ 0 & \dots & \dots & \dots & 0 & 1 & 1 \end{bmatrix}_{n \times n} = \sum_{i=1}^n F_i$$

which is the well-known result from [\[14\]](#).

4. On the Lucas p -numbers and permanent of certain matrices

In the above results we also determine relationships between the permanents of certain square matrices and the Fibonacci p -numbers. Here we determine the similar directions for the Lucas p -numbers. For these purposes, we define a new $(n \times n)$ matrix $H(n, p)$.

Definition 5. For $n > p$ and $p \geq 1$, let $H(n, p) = [h_{ij}]$ be the $n \times n$ matrix with $h_{i+1, i} = h_{i, i+p} = 1$ for a fixed integer p and all $i, j, h_{ii} = 1$ for all i except from $i = p + 1, h_{n-p, n-p} = p + 1$ and 0 otherwise.

Clearly,

$$H(n, p) = \begin{bmatrix} 1 & 0 & \dots & 0 & \overset{(\dot{p} + 1)\text{th}}{\downarrow} 1 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 & 1 & 0 & \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & 0 & 1 & p + 1 & 0 & & & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \ddots & \ddots & \ddots & \dots \\ 0 & 0 & \dots & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & \dots & 0 & 0 & 1 & 1 \end{bmatrix} \rightarrow (n - p)\text{th}$$

Then we have the following [Theorem](#).

Theorem 6. For $n > p \geq 1$,

$$\text{per } H(n, p) = L_p(n).$$

Proof. (Induction on n) First, we consider the case $n = p + 1$. Then the matrix $H(p + 1, p)$ takes the following form:

$$H(p + 1, p) = \begin{bmatrix} p + 1 & 0 & \dots & \dots & 0 & 1 \\ 1 & 1 & 0 & \dots & \dots & 0 \\ 0 & 1 & 1 & 0 & \dots & 0 \\ \vdots & \ddots & \ddots & \ddots & \ddots & \dots \\ 0 & \dots & 0 & 1 & 1 & 0 \\ 0 & \dots & \dots & 0 & 1 & 1 \end{bmatrix}.$$

Thus expanding by the Laplace expansion of permanent with respect to the first row gives us

$$perH(p+1,p) = per \begin{bmatrix} p+1 & 0 & \dots & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \dots \\ \dots & 0 & 1 & 1 & 0 \\ \dots & \dots & 0 & 1 & 1 \end{bmatrix} + per \begin{bmatrix} 1 & 0 & \dots & \dots & 0 \\ 1 & 1 & 0 & \dots & \vdots \\ 0 & 1 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ 0 & \dots & 0 & 1 & 1 \end{bmatrix} = p+2 \tag{14}$$

By the definition of the Lucas p -numbers, we have for $n = p + 1$,

$$L_p(p+1) = L_p(p) + L_p(0) = 1 + p + 1 = p + 2. \tag{15}$$

Considering (14) and (15), we have the conclusion for the first case $n = p + 1$.

Now we consider the case $n > p + 1$. Suppose that the claim is true for $n > p + 1$. Then we show that the claim is true for $n + 1$. Thus if we expand the $perH(n,p)$ by the Laplace expansion of permanent with respect to first row, then we obtain by the definition of matrix $H(n,p)$

$$perH(n+1,p) = perH(n,p) + perH(n-p,p)$$

By our assumption and the definition of Lucas p -numbers, we may write

$$perH(n+1,p) = perH(n,p) + perH(n-p,p) = L_p(n) + L_p(n-p) = L_p(n+1)$$

Thus the proof is complete. \square

For example, when $n = 6$ and $p = 2$, the matrix $H(6,2)$ takes the following form:

$$H(6,2) = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 1 & 0 \\ 0 & 0 & 1 & 3 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

Then by Theorem 6, we have

$$perH(6,2) = 10$$

Indeed by the definition of the Lucas 2-numbers, we see that $L_2(6) = perH(6,2) = 10$.

Define also the $n \times n$ matrix $E(n,p)$ with $e_{i+1,i} = e_{i,i+p} = 1$ for a fixed integer p and all $i, j, e_{ii} = 1$ for all i except from $i = p + 1, e_{p+1,p+1} = p + 1$ and 0 otherwise.

Clearly the matrix $E(n,p)$ have the form

$$E(n,p) = \begin{bmatrix} 1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 & 0 \\ 1 & 1 & 0 & \dots & \dots & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & 1 & 1 & 0 & \dots & 0 & 1 & 0 \\ \dots & 0 & 1 & p+1 & 0 & \dots & 0 & 1 \\ 0 & \dots & 0 & 1 & 1 & 0 & \dots & 0 \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & \dots & \dots & \dots & 0 & 1 & 1 \end{bmatrix}$$

For example when $n = 5, p = 2$, then

$$E(5,2) = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 & 0 \\ 0 & 1 & 3 & 0 & 1 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 1 \end{bmatrix}$$

Considering the definitions of matrices $H(n, p)$ and $E(n, p)$, we have the following corollary without proof by the result of Theorem 6.

Corollary 7. For $n > p \geq 1$

$$\text{per}E(n, p) = L_p(n)$$

Now we derive relationship between the sums of the Lucas p – numbers subscripted from 0 to n and permanent of a certain matrix. For this purpose, we give the following definition.

For compactness, we define the $n \times n$ matrix $G(n, p)$ as in the following form:

$$G(n, p) = \begin{bmatrix} 1 & 1 & \dots & 1 \\ 1 & & & \\ 0 & & E(n-1, p) & \\ \vdots & & & \\ 0 & & & \end{bmatrix}$$

where $E(n, p)$ be as before.

Then we have the following Theorem.

Theorem 8. For $n > p \geq 1$

$$\text{per}G(n, p) = \sum_{k=0}^{n-1} L_p(k)$$

Proof. We will use the induction method to prove Theorem 8. Let $p = 1$ and so $n = 2$. Then

$$\text{per}G(2, 1) = \text{per} \begin{bmatrix} 1 & 1 \\ 1 & 2 \end{bmatrix} = 3$$

Since $L_1(0) = 2, L_1(1) = 1, \sum_{k=0}^1 L_1(k) = 3$. Thus the proof is complete for $n = 2$ and $p = 1$. Consider the case $n = 3$ and $p = 1$. Then

$$\text{per}G(3, 1) = \text{per} \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 0 & 1 & 2 \end{bmatrix} = 6$$

Since also $L_1(2) = 3, \sum_{k=0}^2 L_1(k) = 6$. So the proof is complete for $n = 3$ and $p = 1$. Suppose that the equation holds for n . Then we show that the equation holds for $n + 1$. If we extend the $\text{per}G(n + 1, p)$ according to the first column, then we obtain by the definitions of the matrices $G(n, p)$ and $E(n, p)$

$$\text{per}G(n + 1, p) = \text{per}G(n, p) + E(n, p)$$

By our assumption and the result of Corollary 7, we can write

$$\text{per}G(n + 1, p) = \text{per}G(n, p) + E(n, p) = \sum_{k=0}^{n-1} L_p(k) + L_p(n) = \sum_{k=0}^n L_p(k)$$

Thus the proof is complete. \square

In the above results, we give relationships between the Fibonacci, Lucas p -numbers and the permanents of certain matrices. Here we give relationships between determinants of certain matrices and the Fibonacci and Lucas p -numbers and their sums.

A matrix A is called *convertible* if there is an $n \times n(1, -1)$ -matrix H such that $\text{per}A = \det(A \circ H)$, where $A \circ H$ denotes the Hadamard product of A and H . Such a matrix H is called a *converter* of A .

Let S be a $(1, -1)$ -matrix of order n , defined by

$$S = \begin{bmatrix} 1 & 1 & \dots & 1 & 1 \\ -1 & 1 & \dots & 1 & 1 \\ 1 & -1 & \dots & 1 & 1 \\ \vdots & \vdots & & \vdots & \vdots \\ 1 & 1 & \dots & -1 & 1 \end{bmatrix}$$

Denote $M(n, p) \circ S, T(n, p) \circ S, H(n, p) \circ S$ and $G(n, p) \circ S$ by $\widehat{M}(n, p), \widehat{T}(n, p), \widehat{H}(n, p)$ and $\widehat{G}(n, p)$, respectively. Then we have the following Corollaries without proof.

Corollary 9. For $n > p \geq 1$

$$\det \widehat{M}(n, p) = F_p(n+1)$$

Corollary 10. For $n > p \geq 1$

$$\det \widehat{T}(n, p) = \sum_{i=1}^{n+1} F_p(i)$$

Corollary 11. For $n > p \geq 1$

$$\det \widehat{H}(n, p) = L_p(n)$$

Corollary 12. For $n > p \geq 1$

$$\det \widehat{G}(n, p) = \sum_{k=0}^{n-1} L_p(k).$$

For example, when $n = 5$ and $p = 2$

$$\det \widehat{G}(5, 2) = \begin{vmatrix} 1 & 1 & 1 & 1 & 1 \\ -1 & 1 & 0 & 1 & 0 \\ 0 & -1 & 3 & 0 & 1 \\ 0 & 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & -1 & 1 \end{vmatrix} = 14.$$

Since the definition of the Lucas 2-numbers, we have $\sum_{i=0}^4 L_2(i) = 14$.

References

- [1] Brualdi RA, Gibson PM. Convex polyhedra of Doubly Stochastic matrices I: applications of the Permanent Function. *J Combin Theory A* 1977;22:194–230.
- [2] Byrd PF. Problem B-12: A Lucas determinant. *Fibonacci Quart* 1963;1(4):78.
- [3] Cahill ND, Narayan DA. Fibonacci and Lucas numbers as tridiagonal matrix determinants. *Fibonacci Quart* 2004;42(3):216–21.
- [4] Er MC. Sums of Fibonacci numbers by matrix methods. *Fibonacci Quart* 1984;22(3):204–7.
- [5] Falcon S, Plaza A. On the Fibonacci k -numbers. *Chaos, Solitons & Fractals* 2007;32(5):1615–24.
- [6] Harary F. Determinants, permanents and bipartite graphs. *Math Mag* 1969;42:146–8.
- [7] Kilic E, Tasci D. On families of Bipartite graphs associated with sums of Fibonacci and Lucas numbers. *Ars Combin*, in press.
- [8] Kilic E, Tasci D. On families of Bipartite graphs associated with sums of generalized order- k Fibonacci and Lucas numbers. *Ars Combin*, in press.
- [9] Kilic E, Tasci D. On the second order linear recurrences by tridiagonal matrices. *Ars Combin*, in press.
- [10] Kilic E, Tasci D. On the generalized order – k Fibonacci and Lucas numbers. *Rocky Mountain J Math* 2006;36(6):1915–26.
- [11] Kilic E, Tasci D. On the permanents of some tridiagonal matrices with applications to the Fibonacci and Lucas numbers. *Rocky Mountain. J Math*, in press.
- [12] Kilic E, Tasci D. On the second order linear recurrence satisfied by the permanent of generalized doubly Stochastic matrices. *Ars Combin*, in press.
- [13] Kilic E, Tasci D. On the generalized Fibonacci and Pell sequences by Hessenberg matrices. *Ars Combin*, in press.
- [14] Kilic E, Tasci D. On sums of second order linear recurrences by Hessenberg matrices. *Rocky Mountain J Math*, in press.
- [15] Kilic E, Tasci D, Haukkanen P. On the generalized Lucas sequences by Hessenberg matrices. *Ars Combin*, in press.
- [16] Kilic E. The Binet formula, sums and representations of generalized Fibonacci p -numbers, *European J Combin*, in press, doi:10.1016/j.ejc.2007.03.004.
- [17] Lee G-Y, Lee S-G. A note on generalized Fibonacci numbers. *Fibonacci Quart* 1995;33:273–8.
- [18] Lee G-Y. k -Lucas numbers and associated Bipartite graphs. *Linear Algebra Appl* 2000;320:51–61.

- [19] Lehmer D. Fibonacci and related sequences in periodic tridiagonal matrices. *Fibonacci Quart* 1975;13:150–8.
- [20] Marek-Crnjac L. On the mass spectrum of the elementary particles of the standard model using El Naschie's golden field theory. *Chaos, Solitons & Fractals* 2003;15(4):611–8.
- [21] Marek-Crnjac L. The mass spectrum of high energy elementary particles via El Naschie's golden mean nested oscillators, the Dunkerly–Southwell eigenvalue theorems and KAM. *Chaos, Solitons & Fractals* 2003;18(1):125–33.
- [22] Marek-Crnjac L. The golden mean in the topology of four-manifolds, in conformal field theory, in the mathematical probability theory and in Cantorian space-time. *Chaos, Solitons & Fractals* 2006;28(5):1113–8.
- [23] Maiboroda AO. Finding the golden section in fundamental relations of physical magnitudes, In: Proceedings of the international conference “problems of harmony, symmetry and the golden section in nature, science and art, Vinnitsa, vol. 15; 2003. p. 87–94.
- [24] Mauldin RD, Williams SC. Random recursive construction. *Trans Am Math Soc* 1986;295:325–46.
- [25] Miles Jr EP. Generalized Fibonacci numbers and associated matrices. *Amer Math Monthly* 1960;67:745–52.
- [26] Minc H. Permanents of $(0, 1)$ -Circulants. *Canad Math Bull* 1964;7(2):253–63.
- [27] Minc H. Permanents, *Encyclopedia of mathematics and its applications*. New York: Addison-Wesley; 1978.
- [28] El Naschie MS. On dimension of Cantor set related systems. *Chaos, Solitons & Fractals* 1993;3:675–85.
- [29] El Naschie MS. Quantum mechanics and the possibility of a Cantorian space-time. *Chaos, Solitons & Fractals* 1992;7:485–7.
- [30] El Naschie MS. Is quantum space a random Cantor set with a golden mean dimension at the core? *Chaos, Solitons & Fractals* 1994;4(2):177–9.
- [31] El Naschie MS. Fredholm operators and the wave-particle duality in Cantorian space. *Chaos, Solitons & Fractals* 1998;9(6):975–8.
- [32] El Naschie MS. On a class of general theories for high energy particle physics. *Chaos, Solitons & Fractals* 2002;14(4):649–68.
- [33] El Naschie MS. Complex vacuum fluctuation an chaotic limit” set of any Kleinian group transformation and the mass spectrum of high energy particle physics via spontaneous self-organization. *Chaos, Solitons & Fractals* 2003;17(4):631–8.
- [34] El Naschie MS. On two new fuzzy Kahler manifolds, Klein modular space and “t Hooft holographic principles. *Chaos, Solitons & Fractals* 2006;29:876–81.
- [35] El Naschie MS. From symmetry to particles. *Chaos, Solitons & Fractals* 2007;32:427–30.
- [36] El Naschie MS. SU(5) grand unification in a grand form. *Chaos, Solitons & Fractals* 2007;32:370–4.
- [37] El Naschie MS. Estimation the experimental value of the electromagnetic fine structure constant $\alpha_0 = 1/137.036$ using the Leech lattice in conjunction with the monster group and Spher's kissing number in 24 dimensions. *Chaos, Solitons & Fractals* 2007;32:383–7.
- [38] El Naschie MS. On the topologic ground state of E -infinity spacetime and super string connection. *Chaos, Solitons & Fractals* 2007;32:468–70.
- [39] Petrunenko VV. To the question on physical essence of phenomenon decalogarifmic periodicity, In: Proceedings of the international conference “problems of harmony, symmetry and the golden section in nature, science and art, Vinnitsa, vol. 15; 2003. p. 80–6.
- [40] Stakhov AP. Introduction into algorithmic measurement theory, Soviet Radio, Moscow (1977) [in Russian].
- [41] Stakhov AP. Algorithmic measurement theory, *Znanie, Series “Mathematics and Cybernetics, Moscow (1979) No. 6* [in Russian].
- [42] Stakhov AP. Codes of the golden proportion, Radio and Communications, Moscow (1984) [in Russian].
- [43] Stakhov AP. The golden section in the measurement theory. *Comput Math Appl* 1989;17(4-6):613–38.
- [44] Stakhov AP, Tkachenko IS. Hyperbolic Fibonacci trigonometry. *Rep Ukr Acad Sci* 1993;208(7):9–14 [in Russian].
- [45] Stakhov AP. The golden section and modern harmony mathematics. *Applications of Fibonacci numbers 7*. Kluwer Academic Publishers; 1998. p. 393–99.
- [46] Stakhov AP. A generalization of the Fibonacci Q - matrix. *Rep Natl Acad Sci Ukraine* 1999;9:46–9.
- [47] Stakhov AP. Hyperbolic Fibonacci and Lucas functions: a new mathematics for the living nature, ITI, Vinnitsa (2003).
- [48] Stakhov AP, Rozin B. Theory of Binet formulas for Fibonacci and Lucas p -numbers. *Chaos, Solitons & Fractals* 2006;27:1162–77.
- [49] Stakhov AP. The generalized principle of the golden section and its applications in mathematics, science and engineering. *Chaos, Solitons & Fractals* 2005;26:263–89.
- [50] Stakhov AP. Fibonacci matrices, a generalization of the “Cassini formula”, and a new coding theory, 30(1); 2006: p. 56–66.
- [51] Stakhov AP, Rozin B. The “golden” algebraic equations. *Chaos, Solitons & Fractals* 2006;27:1415–21.
- [52] Stakhov AP. Fundamentals of a new kind of mathematics based on the golden section. *Chaos, Solitons & Fractals* 2005;27:1124–46.
- [53] Stakhov AP. The “golden” matrices and a new kind of cryptography. *Chaos, Solitons & Fractals* 2007;32(3):1138–46.
- [54] Strang G. Introduction to linear algebra. 2nd ed. Wellesley-Cambridge: Wellesley MA; 1998.
- [55] Strang G, Borre K. Linear algebra. Geodesy and GPS. Wellesley-Cambridge: Wellesley MA; 1997. pp. 555–557.
- [56] Tasci D, Kilic E. On the order- k generalized Lucas numbers. *Appl Math Comput* 2004;155(3):637–41.
- [57] Vladimirov YS. *Metaphysic*. Moscow: BINOM: 2002 [in Russian].
- [58] Vorobyov NN, *Fibonacci numbers*, Nauka, Moscow (1978) [in Russian].