

# ON THE SECOND ORDER LINEAR RECURRENCES BY GENERALIZED DOUBLY STOCHASTIC MATRICES

E. KILIC<sup>1</sup> AND D. TASCI<sup>2</sup>

ABSTRACT. In this paper, we consider the relationships between the second order linear recurrences, and the generalized doubly stochastic permanents and determinants.

## 1. INTRODUCTION

The Fibonacci sequence,  $\{F_n\}$ , is defined by the recurrence relation, for  $n \geq 1$

$$F_{n+1} = F_n + F_{n-1} \quad (1.1)$$

where  $F_0 = 0$ ,  $F_1 = 1$ . The Lucas Sequence,  $\{L_n\}$ , is defined by the recurrence relation, for  $n \geq 1$

$$L_{n+1} = L_n + L_{n-1} \quad (1.2)$$

where  $L_0 = 2$ ,  $L_1 = 1$ .

The well-known Fibonacci, Lucas and Pell numbers can be generalized as follows: Let  $A$  and  $B$  be nonzero, relatively prime integers such that  $D = A^2 - 4B \neq 0$ . Define sequences  $\{u_n\}$  and  $\{v_n\}$  by, for all  $n \geq 2$  (see [14]),

$$u_n = Au_{n-1} - Bu_{n-2} \quad (1.3)$$

$$v_n = Av_{n-1} - Bv_{n-2} \quad (1.4)$$

where  $u_0 = 0$ ,  $u_1 = 1$  and  $v_0 = 2$ ,  $v_1 = A$ . If  $A = 1$  and  $B = -1$ , then  $u_n = F_n$  (the  $n$ th Fibonacci number) and  $v_n = L_n$  (the  $n$ th Lucas number). If  $A = 2$  and  $B = -1$ , then  $u_n = P_n$  (the  $n$ th Pell number).

An alternative is to let the roots of the equation  $t^2 - At + B = 0$  be, for  $n \geq 0$

$$u_n = \frac{\sigma^n - \gamma^n}{\sigma - \gamma} \quad \text{and} \quad v_n = \sigma^n + \gamma^n. \quad (1.5)$$

There are many connections between permanents or determinants of tridiagonal matrices and the Fibonacci and Lucas numbers. For example,

---

2000 *Mathematics Subject Classification.* 11B37, 15A15, 15A51.

*Key words and phrases.* Second order linear recurrences, Generalized doubly stochastic matrix, Permanent, Determinant.

Minc [12] define a  $n \times n$  super diagonal  $(0, 1)$  –matrix  $F(n, k)$  for  $n > k \geq 2$ , and show that the permanent of  $F(n, k)$  equals to the generalized order- $k$  Fibonacci numbers. Also he give some relations involving the permanents of some  $(0, 1)$  –Circulant matrices and the usual Fibonacci numbers.

In [8], the authors present a nice result involving the permanent of an  $(-1, 0, 1)$ -matrix and the Fibonacci Number  $F_{n+1}$ . The authors then explore similar directions involving the positive subscripted Fibonacci and Lucas Numbers as well as their uncommon negatively subscripted counterparts. Finally the authors explore the generalized order- $k$  Lucas numbers, (see [19] and [7] for more detail the generalized Fibonacci and Lucas numbers), and their permanents.

In [9] and [10], the authors gave the relations involving the generalized Fibonacci and Lucas numbers and the permanent of the  $(0, 1)$  –matrices. The results of Minc, [12], and the result of Lee, [9], on the generalized Fibonacci numbers are the same because they use the same matrix. However, Lee proved the same result by a different method, contraction method for the permanent (for more detail of the contraction method see [1]).

In [11], Lehmer proves a very general result on permanents of tridiagonal matrices whose main diagonal and super-diagonal elements are ones and whose subdiagonal entries are somewhat arbitrary.

Also in [16] and [17], the authors define a family of tridiagonal matrices  $M(n)$  and show that the determinants of  $M(n)$  are the Fibonacci numbers  $F_{2n+2}$ . In [4] and [3], the family of tridiagonal matrices  $H(n)$  and the authors show that the determinants of  $H(n)$  are the Fibonacci numbers  $F_n$ . In a similar family of matrices, the  $(1, 1)$  element of  $H(n)$  is replaced with a 3. The determinants, [2], now generate the Lucas sequence  $L_n$ .

In [5], the authors find the families of  $(0, 1)$  –matrices such that permanents of the matrices, equal to the sums of Fibonacci and Lucas numbers.

Recently, in [6], the authors define two tridiagonal matrices and then give the relationships the permanents and determinants of these matrices and the second order linear recurrences.

The *permanent* of an  $n$ -square matrix  $A = (a_{ij})$  is defined by

$$\text{per}A = \sum_{\sigma \in S_n} \prod_{i=1}^n a_{i\sigma(i)}$$

where the summation extends over all permutations  $\sigma$  of the symmetric group  $S_n$ . Also one can find more applications of permanents in [13].

Let  $A = [a_{ij}]$  be an  $m \times n$  real matrix row vectors  $\alpha_1, \alpha_2, \dots, \alpha_m$ . We say  $A$  is *contractible on column* (resp. *row*)  $k$  if *column* (resp. *row*)  $k$  contains exactly two nonzero entries. Suppose  $A$  is contractible on column  $k$  with  $a_{ik} \neq 0 \neq a_{jk}$  and  $i \neq j$ . Then the  $(m-1) \times (n-1)$  matrix  $A_{ij:k}$  obtained from  $A$  by replacing row  $i$  with  $a_{jk}\alpha_i + a_{ik}\alpha_j$  and deleting row

$j$  and column  $k$  is called *the contraction of  $A$  on column  $k$  relative to rows  $i$  and  $j$* . If  $A$  is contractible on row  $k$  with  $a_{ki} \neq 0 \neq a_{kj}$  and  $i \neq j$ , then the matrix  $A_{k:i,j} = \left[ A_{i,j;k}^T \right]^T$  is called *the contraction of  $A$  on row  $k$  relative to columns  $i$  and  $j$* . Every contraction used in this paper will be on the first column using the first and second rows. We say that  $A$  can be contracted to a matrix  $B$  if either  $B = A$  or exist matrices  $A_0, A_1, \dots, A_t$  ( $t \geq 1$ ) such that  $A_0 = A$ ,  $A_t = B$ , and  $A_r$  is a contraction of  $A_{r-1}$  for  $r = 1, 2, \dots, t$ . One can find the following fact in [1]: let  $A$  be a nonnegative integral matrix of order  $n > 1$  and let  $B$  be a contraction of  $A$ . Then

$$\text{per } A = \text{per } B. \quad (1.6)$$

We also recall the following definitions :

**Definition 1.** A matrix  $A = (a_{ij})$  of order  $n$  is said to be nonnegative if  $a_{ij} \geq 0$ ,  $i, j = 1, 2, \dots, n$ .

**Definition 2.** A nonnegative  $n \times n$  matrix  $A$  is called row stochastic, or simply stochastic, if all its rows sum 1.

**Definition 3.** A nonnegative  $n \times n$  matrix  $A$  is called row stochastic, if all its rows and columns sum 1.

We give the following definitions (see [15] and [18], respectively).

**Definition 4.** A matrix  $A = (a_{kj})$  of order  $n$  is said to be generalized stochastic if

$$\sum_{j=1}^n a_{kj} = s, \quad k = 1, 2, \dots, n$$

where  $s$  is a complex number.

**Definition 5.** If  $A = (a_{kj})$  is such that

$$\sum_{j=1}^n a_{kj} = s, \quad k = 1, 2, \dots, n \quad \text{and} \quad \sum_{k=1}^n a_{kj} = s, \quad j = 1, 2, \dots, n$$

then  $A$  is said to be generalized doubly stochastic matrix.

Note that a generalized stochastic or generalized doubly stochastic matrix need to be nonnegative.

In this paper, we give the relationships between the permanents of some generalized symmetric doubly stochastic matrices and the second order linear recurrences.

## 2. THE MAIN RESULTS

In this section, we define a  $n \times n$  generalized symmetric doubly stochastic matrix  $D_n$  and then show that its permanent equals to the  $n$ th term of the sequence  $\{v_n\}$ .

We define a  $n \times n$  generalized symmetric doubly stochastic matrix  $D_n$  with  $d_{11} = \frac{\alpha}{\alpha+\beta}$ ,  $d_{ii} = 0$  for  $2 \leq i \leq n-1$ , let  $n$  be an even number,  $d_{2k,2k+1} = \frac{\alpha}{\alpha+\beta}$  for  $1 \leq k \leq \frac{n-2}{2}$ ,  $d_{2k-1,2k} = \frac{\beta}{\alpha+\beta}$  for  $1 \leq k \leq \frac{n}{2}$  and  $d_{nn} = \frac{\alpha}{\alpha+\beta}$ , and, let  $n$  be an odd number,  $d_{2k,2k+1} = \frac{\alpha}{\alpha+\beta}$  and  $d_{2k-1,2k} = \frac{\beta}{\alpha+\beta}$  for  $1 \leq k \leq \frac{n-1}{2}$ , and  $d_{nn} = \frac{\beta}{\alpha+\beta}$ . Clearly, if  $n$  is an even, then

$$D_n = \begin{bmatrix} \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} & & & & & & 0 \\ \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} & & & & & \\ & \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} & & & & \\ & & \frac{\beta}{\alpha+\beta} & \ddots & \ddots & & & \\ & & & \ddots & 0 & \frac{\beta}{\alpha+\beta} & & \\ 0 & & & & \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} & & \end{bmatrix}.$$

We note that the rows and columns sums of the matrix  $D_n$  equal to 1. However, in general, since the entries of the matrix  $D_n$ ,  $\frac{\alpha}{\alpha+\beta}$  and  $\frac{\beta}{\alpha+\beta}$  are not nonnegative, the matrix  $D_n$  is a generalized doubly stochastic matrix.

Then we have the following Theorem.

**Theorem 1.** *Let the matrix  $D_n$  be as before. Then, for  $n \geq 2$*

$$\text{per}D_n = \frac{\alpha^n + \beta^n}{(\alpha + \beta)^n}.$$

*Proof.* If  $n = 2$ , then we have

$$\text{per}D_2 = \text{per} \begin{bmatrix} \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} \\ \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix} = \frac{\alpha^2 + \beta^2}{(\alpha + \beta)^2}.$$

If  $n = 3$ , then we have

$$\text{per}D_3 = \text{per} \begin{bmatrix} \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} & \\ \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} \\ & \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} \end{bmatrix} = \frac{\alpha^3 + \beta^3}{(\alpha + \beta)^3}.$$

We suppose that  $n$  is an even number and let  $D_n^k$  be the  $k$ th contraction of  $D_n$ ,  $1 \leq k \leq n-2$ . Since the definition of the matrix  $D_n$ , the matrix  $D_n$  can be contracted on column 1 so that

$$D_n^1 = \begin{bmatrix} \frac{\beta^2}{(\alpha+\beta)^2} & \frac{\alpha^2}{(\alpha+\beta)^2} & & & & 0 \\ \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} & & & \\ & \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} & & \\ & & \frac{\alpha}{\alpha+\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{\beta}{\alpha+\beta} \\ 0 & & & & \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix}.$$

Since the matrix  $D_n^1$  can be contracted on column 1,

$$D_n^2 = \begin{bmatrix} \frac{\alpha^3}{(\alpha+\beta)^3} & \frac{\beta^3}{(\alpha+\beta)^3} & & & & 0 \\ \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} & & & \\ & \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} & & \\ & & \frac{\beta}{\alpha+\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{\beta}{\alpha+\beta} \\ 0 & & & & \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix}.$$

Continuing this process, we have, for even number  $k$ ,

$$D_n^k = \begin{bmatrix} \frac{\alpha^{k+1}}{(\alpha+\beta)^{k+1}} & \frac{\beta^{k+1}}{(\alpha+\beta)^{k+1}} & & & & 0 \\ \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} & & & \\ & \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} & & \\ & & \frac{\beta}{\alpha+\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{\beta}{\alpha+\beta} \\ 0 & & & & \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix}$$

for  $3 \leq k \leq n-4$ . Hence,

$$D_n^{(n-3)} = \begin{bmatrix} \frac{\beta^{n-2}}{(\alpha+\beta)^{n-2}} & \frac{\alpha^{n-2}}{(\alpha+\beta)^{n-2}} & 0 \\ \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} \\ 0 & \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix}$$

which, by contraction of  $D_n^{(n-4)}$  on column 1, gives

$$D_n^{(n-2)} = \begin{bmatrix} \frac{\alpha^{n-1}}{(\alpha+\beta)^{n-1}} & \frac{\beta^{n-1}}{(\alpha+\beta)^{n-1}} \\ \frac{\beta}{\alpha+\beta} & \frac{\alpha}{\alpha+\beta} \end{bmatrix}.$$

By applying (1.6), we have  $\text{per} D_n = \text{per} D_n^{(n-2)} = \frac{\alpha^n + \beta^n}{(\alpha+\beta)^n}$ .

When the case  $n$  is odd number, the proof easily follows from the above case  $n$  is even number.

So the proof is complete.  $\square$

Now we give a relationship between the result of Theorem 1 and second order recurrence  $\{v_n\}$ .

**Corollary 1.** *Let  $\alpha$  and  $\beta$  be the roots of the equation  $t^2 - At + B = 0$ . Then, for  $n \geq 2$*

$$\text{per} D_n = \frac{v_n}{A^n}$$

where  $v_n$  is the  $n$ th term of the sequence  $\{v_n\}$  and  $A = \alpha + \beta$ .

*Proof.* For the sequence  $\{v_n\}$ , the Binet formula is given by  $v_n = \alpha^n + \beta^n$  and since Theorem 1, the proof is easily seen.  $\square$

For example, when  $\alpha = \frac{1+\sqrt{5}}{2}$  and  $\beta = \frac{1-\sqrt{5}}{2}$ , by Corollary 1, we have

$$\text{per} \begin{bmatrix} \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} & & & & 0 \\ \frac{\beta}{\alpha+\beta} & 0 & \frac{\alpha}{\alpha+\beta} & & & \\ & \frac{\alpha}{\alpha+\beta} & 0 & \frac{\beta}{\alpha+\beta} & & \\ & & \frac{\beta}{\alpha+\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{\alpha}{\alpha+\beta} \\ 0 & & & & \frac{\alpha}{\alpha+\beta} & \frac{\beta}{\alpha+\beta} \end{bmatrix}_{n \times n} = L_n$$

where  $L_n$  is the  $n$ th Lucas number.

Second, we define a  $n \times n$  generalized symmetric doubly stochastic matrix  $H_n$  with  $h_{11} = \frac{\alpha}{\alpha-\beta}$ ,  $h_{ii} = 0$  for  $2 \leq i \leq n-1$ , let  $n$  be an even number,  $h_{2k,2k+1} = \frac{\alpha}{\alpha-\beta}$  for  $1 \leq k \leq \frac{n-2}{2}$ ,  $h_{2k-1,2k} = \frac{-\beta}{\alpha-\beta}$  for  $1 \leq k \leq \frac{n}{2}$  and  $h_{nn} = \frac{\alpha}{\alpha-\beta}$ , and, let  $n$  be an odd number,  $h_{2k,2k+1} = \frac{\alpha}{\alpha-\beta}$  and  $h_{2k-1,2k} = \frac{-\beta}{\alpha-\beta}$  for  $1 \leq k \leq \frac{n-1}{2}$ , and  $h_{nn} = \frac{-\beta}{\alpha-\beta}$ . Clearly, for even number  $n$ , the matrix

$$H_n = \begin{bmatrix} \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} & & & & 0 \\ \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & \\ & \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & \\ & & \frac{-\beta}{\alpha-\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{-\beta}{\alpha-\beta} \\ 0 & & & & \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix}.$$

We note that the rows and columns sums of the matrix  $H_n$  equal to 1. However, in general, since the entries of the matrix  $H_n$ ,  $\frac{\alpha}{\alpha-\beta}$  and  $\frac{-\beta}{\alpha-\beta}$  are not nonnegative, the matrix  $H_n$  is a generalized doubly stochastic matrix.

Then we have the following Theorem.

**Theorem 2.** *Let the generalized doubly stochastic matrix  $H_n$  be as before. Then, for  $n \geq 2$*

$$\text{per}H_n = \begin{cases} \frac{\alpha^n - \beta^n}{(\alpha - \beta)^n} & \text{if } n \text{ is odd number,} \\ \frac{\alpha^n + \beta^n}{(\alpha - \beta)^n} & \text{if } n \text{ is even number.} \end{cases}$$

*Proof.* We consider the first case  $n$  is odd number. If  $n = 3$ , then we have

$$\text{per}H_3 = \begin{bmatrix} \frac{\alpha}{\alpha - \beta} & \frac{-\beta}{\alpha - \beta} & 0 \\ \frac{-\beta}{\alpha - \beta} & 0 & \frac{\alpha}{\alpha - \beta} \\ 0 & \frac{\alpha}{\alpha - \beta} & -\beta \end{bmatrix} = \frac{\alpha^3 - \beta^3}{(\alpha - \beta)^3}.$$

If  $n = 5$ , then we have

$$\text{per}H_5 = \begin{bmatrix} \frac{\alpha}{\alpha - \beta} & \frac{-\beta}{\alpha - \beta} & & & \\ \frac{-\beta}{\alpha - \beta} & 0 & \frac{\alpha}{\alpha - \beta} & & \\ & \frac{\alpha}{\alpha - \beta} & 0 & \frac{-\beta}{\alpha - \beta} & \\ & & \frac{-\beta}{\alpha - \beta} & 0 & \frac{\alpha}{\alpha - \beta} \\ & & & \frac{\alpha}{\alpha - \beta} & \frac{-\beta}{\alpha - \beta} \end{bmatrix} = \frac{\alpha^5 - \beta^5}{(\alpha - \beta)^5}.$$

Let  $H_n^k$  be the  $k$ th contraction of  $H_n$ ,  $1 \leq k \leq n - 2$ . Since the definition of the matrix  $H_n$ , the matrix  $H_n$  can be contracted on column 1 so that

$$H_n^1 = \begin{bmatrix} \frac{\beta^2}{(\alpha - \beta)^2} & \frac{\alpha^2}{(\alpha - \beta)^2} & & & \\ \frac{\alpha}{\alpha - \beta} & 0 & \frac{-\beta}{\alpha - \beta} & & \\ & \frac{-\beta}{\alpha - \beta} & 0 & \frac{\alpha}{\alpha - \beta} & \\ & & \frac{\alpha}{\alpha - \beta} & \ddots & \ddots \\ & & & \ddots & 0 & \frac{\alpha}{\alpha - \beta} \\ & & & & \frac{\alpha}{\alpha - \beta} & \frac{-\beta}{\alpha - \beta} \end{bmatrix}.$$



Since the matrix  $H_n^1$  can be contracted on column 1,

$$H_n^2 = \begin{bmatrix} \frac{\alpha^3}{(\alpha-\beta)^3} & \frac{-\beta^3}{(\alpha-\beta)^3} & & & & & & & \\ \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & & & & \\ & \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & & & & \\ & & & \frac{-\beta}{\alpha-\beta} & \ddots & \ddots & & & \\ & & & & \ddots & & 0 & \frac{\alpha}{\alpha-\beta} & \\ & & & & & & & \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \end{bmatrix}.$$

Continuing this process, we have, for odd number  $k$ ,

$$H_n^k = \begin{bmatrix} \frac{\beta^{k+1}}{(\alpha-\beta)^{k+1}} & \frac{\alpha^{k+1}}{(\alpha-\beta)^{k+1}} & & & & & & & \\ \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & & & & & \\ & \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & & & \\ & & & \frac{\alpha}{\alpha-\beta} & \ddots & \ddots & & & \\ & & & & \ddots & & 0 & \frac{\alpha}{\alpha-\beta} & \\ & & & & & & & \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \end{bmatrix}$$

for  $3 \leq k \leq n-4$ . Hence,

$$H_n^{(n-3)} = \begin{bmatrix} \frac{\alpha^{n-2}}{(\alpha-\beta)^{n-2}} & \frac{-\beta^{n-2}}{(\alpha-\beta)^{n-2}} & 0 \\ \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} \\ 0 & \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \end{bmatrix}$$

which, by contraction of  $H_n^{(n-4)}$  on column 1, gives

$$H_n^{(n-2)} = \begin{bmatrix} \frac{\beta^{n-1}}{(\alpha-\beta)^{n-1}} & \frac{\alpha^{n-1}}{(\alpha-\beta)^{n-1}} \\ \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \end{bmatrix}.$$

By applying (1.6), we have  $Per H_n = per H_n^{(n-2)} = \frac{\alpha^n - \beta^n}{(\alpha-\beta)^n}$ .

Now we consider the second case  $n$  is even number. If  $n = 2$ , then we have

$$\text{per}H_2 = \begin{bmatrix} \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \\ \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix} = \frac{\alpha^2 + \beta^2}{(\alpha - \beta)^2}.$$

Taking successive contractions of the matrix  $H_n$ , similarly the above, we have, for even number  $k$ ,

$$H_n^k = \begin{bmatrix} \frac{\alpha^{k+1}}{(\alpha-\beta)^{k+1}} & \frac{-\beta^{k+1}}{(\alpha-\beta)^{k+1}} & & & & & \\ & \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & \\ & & \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & \\ & & & \frac{-\beta}{\alpha-\beta} & \ddots & \ddots & \\ & & & & \ddots & 0 & \frac{-\beta}{\alpha-\beta} \\ & & & & & \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix}$$

for  $3 \leq k \leq n - 4$ . Hence,

$$H_n^{(n-3)} = \begin{bmatrix} \frac{\beta^{n-2}}{(\alpha-\beta)^{n-2}} & \frac{\alpha^{n-2}}{(\alpha-\beta)^{n-2}} & 0 \\ \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} \\ 0 & \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix}$$

which, by contraction of  $H_n^{(n-4)}$  on column 1, gives

$$H_n^{(n-2)} = \begin{bmatrix} \frac{\alpha^{n-1}}{(\alpha-\beta)^{n-1}} & \frac{-\beta^{n-1}}{(\alpha-\beta)^{n-1}} \\ \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix} ..$$

By applying (1.6), we have  $\text{Per}H_n = \text{per}H_n^{(n-2)} = \frac{\alpha^n + \beta^n}{(\alpha - \beta)^n}$  for even number  $n$ .

So the proof is complete.  $\square$

Now we give a relationship between the result of Theorem 2 and sequence  $\{u_n\}$ .

**Corollary 2.** Let  $\alpha$  and  $\beta$  be the roots of the equation  $t^2 - At + B = 0$ . Then, for  $n > 1$

$$\text{per}H_n = \begin{cases} \frac{u_n}{(\alpha-\beta)^{n-1}} & \text{if } n \text{ is odd number,} \\ \frac{v_n}{(\alpha-\beta)^n} & \text{if } n \text{ is even number.} \end{cases}$$

where  $u_n$  and  $v_n$  are the  $n$ th terms of the sequences  $\{u_n\}$  and  $\{v_n\}$ .

*Proof.* Considering the Binet formulas for the sequences  $\{u_n\}$  and  $\{v_n\}$ , and the result of Theorem 2, the proof is easily seen.  $\square$

For example, when  $\alpha = \frac{1+\sqrt{5}}{2}$  and  $\beta = \frac{1-\sqrt{5}}{2}$ , by Corollary 2, we have

$$\text{per} \begin{bmatrix} \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} & & & & 0 \\ \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & \\ & \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & \\ & & \frac{-\beta}{\alpha-\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{\alpha}{\alpha-\beta} \\ 0 & & & & \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} \end{bmatrix}_{n \times n} = \frac{F_n}{(\sqrt{5})^{n-1}}$$

for even number  $n$ , where  $F_n$  is the  $n$ th Fibonacci number, and

$$\text{per} \begin{bmatrix} \frac{\alpha}{\alpha-\beta} & \frac{-\beta}{\alpha-\beta} & & & & 0 \\ \frac{-\beta}{\alpha-\beta} & 0 & \frac{\alpha}{\alpha-\beta} & & & \\ & \frac{\alpha}{\alpha-\beta} & 0 & \frac{-\beta}{\alpha-\beta} & & \\ & & \frac{-\beta}{\alpha-\beta} & \ddots & \ddots & \\ & & & \ddots & 0 & \frac{-\beta}{\alpha-\beta} \\ 0 & & & & \frac{-\beta}{\alpha-\beta} & \frac{\alpha}{\alpha-\beta} \end{bmatrix} = \frac{L_n}{(\sqrt{5})^n}$$

for odd number  $n$ , where  $L_n$  is the  $n$ th Lucas number.

#### REFERENCES

- [1] R. A. Brualdi & P. M. Gibson. "Convex polyhedra of Doubly Stochastic matrices I: Applications of the permanent function." *J. Combin. Theory A* **22** (1977): 194-230.

- [2] P.F. Byrd. "Problem B-12: A Lucas determinant." *Fib. Quart.* **1.4** (1963): 78.
- [3] N. D. Cahill and D. A. Narayan. "Fibonacci and Lucas numbers as tridiagonal matrix determinants." *Fib. Quart.* **42.3** (2004): 216-221.
- [4] N. D. Cahill, J. R. D'Errica, D. A. Narayan and J. Y. Narayan. "Fibonacci Determinants." *College math. J.* **3.3** (2002): 221-225.
- [5] E. Kilic and D. Tasci. "On families of bipartite graphs associated with sums of Fibonacci and Lucas numbers." *Ars Combin.* (to appear).
- [6] E. Kilic and D. Tasci, "On the second order linear recurrences by tridiagonal matrices." *Ars Combin.* (to appear).
- [7] E. Kilic and D. Tasci. "On the generalized order- $k$  Fibonacci and Lucas numbers." *Rocky Mountain J. Math.* **36.6** (2006): 1915-1926.
- [8] E. Kilic and D. Tasci, "On the permanents of some tridiagonal matrices with applications to the Fibonacci and Lucas numbers." *Rocky Mountain J. Math.* **37.6** (2007): 1953-1969.
- [9] G. -Y. Lee and S. -G. Lee. "A note on generalized Fibonacci numbers." *Fib. Quart.* **33** (1995): 273-278.
- [10] G. -Y. Lee. " $k$ - Lucas numbers and associated Bipartite graphs." *Linear Algebra Appl.* **320** (2000): 51-61.
- [11] D. Lehmer. "Fibonacci and related sequences in periodic tridiagonal matrices." *Fib. Quart.* **13** (1975): 150-158.
- [12] H. Minc. "Permanents of (0,1)-Circulants." *Canad. Math. Bull.* **7.2** (1964): 253-263.
- [13] H. Minc. *Permanents, Encyclopedia of Mathematics and its Applications.* Addison-Wesley, New York, 1978.
- [14] N. Robbins. *Beginning Number Theory.* Dubuque, Iowa : Wm. C. Brown Publishers 1993.
- [15] F. Salzmann, A note on eigenvalues of nonnegative matrices, *Linear Alg. Appl.* 5, 1972: 329-338.
- [16] G. Strang. *Introduction to Linear Algebra.* 2nd Edition, Wellesley MA, Wellesley-Cambridge, 1998.
- [17] G. Strang and K. Borre. *Linear Algebra. Geodesy and GPS.* Wellesley MA, Wellesley-Cambridge, 1997, pp. 555-557.
- [18] R. L. Soto, The Inverse Spectrum Problem for Positive Generalized Stochastic Matrices,
- [19] D. Tasci and E. Kilic, "On the order- $k$  generalized Lucas numbers." *Appl. Math. Comput.* **155.3** (2004), 637-641.

TOBB ECONOMICS AND TECHNOLOGY UNIVERSITY MATHEMATICS DEPARTMENT 06560  
ANKARA TURKEY

*E-mail address:* [ekilic@etu.edu.tr](mailto:ekilic@etu.edu.tr)

<sup>2</sup>GAZI UNIVERSITY, MATHEMATICS DEPARTMENT, 06500 ANKARA TURKEY