

SOME DOUBLE BINOMIAL SUMS RELATED WITH THE FIBONACCI, PELL AND GENERALIZED ORDER- k FIBONACCI NUMBERS

EMRAH KILIÇ AND HELMUT PRODINGER

ABSTRACT. We consider some double binomial sums related with the Fibonacci, Pell numbers and a multiple binomial sums related with the generalized order- k Fibonacci numbers. The Lagrange-Bürmann formula and other known techniques are used to prove them.

1. INTRODUCTION

The generating function of the Fibonacci numbers F_n is

$$\sum_{n=0}^{\infty} F_n x^n = \frac{x}{1 - x - x^2}.$$

Similarly, the generating function of the Pell numbers P_n is

$$\sum_{n=0}^{\infty} P_n x^n = \frac{x}{1 - 2x - x^2}.$$

The generalized order- k Fibonacci numbers $f_n^{(k)}$ are defined by

$$f_n^{(k)} = \sum_{i=1}^k f_{n-i}^{(k)} \quad \text{for } n > k$$

with initial conditions $f_j^{(k)} = 2^{j-1}$ for $1 \leq j \leq k$.

For example, when $k = 3$, the generalized Fibonacci numbers $f_n^{(3)}$ are reduced to the Tribonacci numbers T_n defined by

$$T_n = T_{n-1} + T_{n-2} + T_{n-3}$$

with $T_1 = 1$, $T_2 = 2$ and $T_3 = 4$, for $n > 3$.

For these number sequences, we recall the combinatorial representations due to [2, 3, 5]:

$$\sum_{i=1}^n \binom{n-i}{i-1} = F_n, \tag{1.1}$$

$$\sum_{i=0}^{\lfloor (n-1)/2 \rfloor} \binom{n}{2i+1} 2^i = P_n, \tag{1.2}$$

2000 *Mathematics Subject Classification.* Primary 05C38, 15A15; Secondary 05A15, 15A18.
Key words and phrases. Fibonacci numbers, generating functions, Lagrange-Bürmann formula.

$$\sum_{i=0}^n \sum_{j=0}^n \binom{n-i}{j} \binom{n-j}{i} = F_{2n+3}. \quad (1.3)$$

Among the formulas (1.1–1.3), the last formula seems to be different from first two identities just above since it includes double sums, see [2]. The authors of the above cited papers use a combinatorial approach to prove these results. For many similar identities, we refer to [6].

In this paper, we shall derive some new double binomial sums related with the Fibonacci, Pell and generalized order- k Fibonacci numbers and then use the Lagrange-Bürmann formula and well known other techniques to prove them.

The Lagrange-Bürmann formula is a very useful tool if one knows a series expansion for $y(x)$ but would like to obtain the series for x in terms of y . We recall the formula (for details see [1, 4]): Suppose a series for y in powers of x is required when $y = x\Phi(y)$. Assume that Φ is analytic in a neighborhood of $y = 0$ with $\Phi(0) \neq 0$. Then

$$x = y/\Phi(y) = \sum_{n=1}^{\infty} a_n y^n, \quad a_1 \neq 0.$$

Then the two (equivalent) versions of the Lagrange(-Bürmann) inversion formula can be written as

$$F(y) = F(0) + \sum_{n=1}^{\infty} \frac{x^n}{n!} \left[\frac{d^{n-1}}{dy^{n-1}} (F'(y)\Phi^n(y)) \right]_{x=0}$$

or

$$\frac{F(y)}{1 - x\Phi'(y)} = \sum_{n=0}^{\infty} \frac{x^n}{n!} \left[\frac{d^n}{dy^n} (F(y)\Phi^n(y)) \right]_{x=0}.$$

We would like to rephrase this using the notation of the “coefficient-of” operator:

$$\frac{F(y)}{1 - x\Phi'(y)} = \sum_{n=0}^{\infty} [y^n] (F(y)\Phi^n(y)) \cdot x^n;$$

we will use it in this form.

2. DOUBLE BINOMIAL SUMS

We start with a result related to Fibonacci numbers:

Theorem 1. For $n > 0$,

$$F_{4n-1} = \sum_{0 \leq i, j \leq n} \binom{n+i}{2j} \binom{n+j}{2i}.$$

Proof. We start from

$$[y^{2j}](1+y)^{n+i} = \binom{n+i}{2j}$$

and compute

$$\begin{aligned} S &= \sum_{i=0}^n (1+y)^{n+i} \binom{n+j}{2i} \\ &= \sum_{i \geq 0} (1+y)^{n+i/2} \binom{n+j}{i} \frac{1+(-1)^i}{2} \end{aligned}$$

$$= \left[\left(1 + \sqrt{1+y}\right)^{j+n} + \left(1 - \sqrt{1+y}\right)^{j+n} \right] \frac{(1+y)^n}{2},$$

here the desired sum takes the form:

$$\begin{aligned} & \sum_{j=0}^n [y^{2j}] \left[\left(1 + \sqrt{1+y}\right)^{j+n} + \left(1 - \sqrt{1+y}\right)^{j+n} \right] \frac{(1+y)^n}{2} \\ &= \sum_{j \geq 0} [y^{2j}] \left(1 + \sqrt{1+y}\right)^{j+n} \frac{(1+y)^n}{2} + \sum_{j \geq 0} [y^{2j}] \left(1 - \sqrt{1+y}\right)^{j+n} \frac{(1+y)^n}{2} \\ &= \sum_{j \geq 0} [y^j] \left(1 + \sqrt{1+y}\right)^{j/2+n} \frac{(1+y)^n}{2} \frac{1 + (-1)^j}{2} + \sum_{j \geq 0} [y^{2j}] \left(1 - \sqrt{1+y}\right)^{j+n} \frac{(1+y)^n}{2}. \end{aligned}$$

Let us consider the first sum:

$$\sum_{j \geq 0} [y^j] \left(1 + \sqrt{1+y}\right)^{j/2+n} (1+y)^n.$$

This is of the form

$$\sum_{j \geq 0} [y^j] F(y) \Phi(y)^j$$

with

$$F(y) = \left(1 + \sqrt{1+y}\right)^n (1+y)^n \quad \text{and} \quad \Phi(y) = \sqrt{1 + \sqrt{1+y}}.$$

The Lagrange-Bürmann formula can now be applied to this sum. The general formula is given by

$$\sum_{j \geq 0} [y^j] F(y) \Phi(y)^j \cdot x^j = \frac{F(y)}{1 - x\Phi'(y)}.$$

We need the instance $x = 1$ here, and the variables x and y are linked via $y = x\Phi(y)$. Notice that $\Phi(y)$ must be a power series in y with a constant term different from zero. Therefore by the solution of $y = \Phi(y)$, we find $y = \alpha = (1 + \sqrt{5})/2$ and so

$$y = \frac{1 + \sqrt{5}}{2}, \quad F(\alpha) = \left(\frac{7 + 3\sqrt{5}}{2} \right)^n,$$

$$\Phi'(\alpha) = \frac{3 - \sqrt{5}}{8}, \quad \frac{1}{1 - \Phi'(\alpha)} = 2 \left(1 - \frac{1}{\sqrt{5}} \right).$$

So our evaluation is

$$2 \left(1 - \frac{1}{\sqrt{5}} \right) \left(\frac{7 + 3\sqrt{5}}{2} \right)^n.$$

The second term is

$$\sum_{j \geq 0} [y^j] \left(1 + \sqrt{1+y}\right)^{j/2+n} (1+y)^n (-1)^j.$$

This is the instance $x = -1$, which translates to $y = -1$ and so the second term is

$$\frac{F(-1)}{1 + \Phi'(-1)} = 0.$$

The last sum is

$$\begin{aligned} \sum_{j \geq 0} [y^{2j}] \left(1 - \sqrt{1+y}\right)^{j+n} (1+y)^n &= \sum_{j \geq 0} [y^{2j}] y^{j+n} \left(\frac{1 - \sqrt{1+y}}{y}\right)^{j+n} (1+y)^n \\ &= \sum_{j \geq 0} [y^j] y^n \left(\frac{1 - \sqrt{1+y}}{y}\right)^{j+n} (1+y)^n. \end{aligned}$$

This is again of the form

$$\sum_{j \geq 0} [y^j] F(y) \Phi(y)^j$$

with

$$F(y) = \left(1 - \sqrt{1+y}\right)^n (1+y)^n \quad \text{and} \quad \Phi(y) = \frac{1 - \sqrt{1+y}}{y}.$$

We need the instance $x = 1$ here, and the link is

$$y = x \left(\frac{1 - \sqrt{1+y}}{y}\right).$$

By the solution of the last equation, we find $y = \beta$ where $\beta = (1 - \sqrt{5})/2$ and so we write

$$y = \beta = \frac{1 - \sqrt{5}}{2}, \quad F(\beta) = \left(\frac{7 - 3\sqrt{5}}{2}\right)^n \quad \text{and} \quad \frac{1}{1 - \Phi'(\alpha)} = 1 + \frac{1}{\sqrt{5}}.$$

So our evaluation is

$$\left(1 + \frac{1}{\sqrt{5}}\right) \left(\frac{7 - 3\sqrt{5}}{2}\right)^n.$$

Altogether

$$\left[\left(1 - \frac{1}{\sqrt{5}}\right) \left(\frac{7 + 3\sqrt{5}}{2}\right)^n + \left(1 + \frac{1}{\sqrt{5}}\right) \left(\frac{7 - 3\sqrt{5}}{2}\right)^n \right] \frac{1}{2} = \frac{\alpha^{4n-1} - \beta^{4n-1}}{\sqrt{5}} = F_{4n-1},$$

as desired. \square

Theorem 2. For $n > 0$,

$$F_{4n+1} = \sum_{1 \leq i, j \leq n+1} \binom{n+i}{2j-1} \binom{n+j}{2i-1}.$$

Proof. Since

$$[y^{2j-1}] (1+y)^{n+i} = \binom{n+i}{2j-1}$$

and

$$\begin{aligned} S &= \sum_{i=1}^{n+1} (1+y)^{n+i} \binom{n+j}{2i-1} \\ &= \sum_{i \geq 0} (1+y)^{n+(i+1)/2} \binom{n+j}{i} \frac{1 - (-1)^i}{2} \\ &= \left[\left(1 + \sqrt{1+y}\right)^{j+n} - \left(1 - \sqrt{1+y}\right)^{j+n} \right] \frac{(1+y)^{n+1/2}}{2}, \end{aligned}$$

here the desired sum takes the form:

$$\begin{aligned}
& \sum_{j=1}^{n+1} [y^{2j-1}] \left[(1 + \sqrt{1+y})^{j+n} - (1 - \sqrt{1+y})^{j+n} \right] \frac{(1+y)^{n+1/2}}{2} \\
&= \sum_{j \geq 1} [y^{2j-1}] (1 + \sqrt{1+y})^{j+n} \frac{(1+y)^{n+1/2}}{2} \\
&\quad - \sum_{j \geq 1} [y^{2j-1}] (1 - \sqrt{1+y})^{j+n} \frac{(1+y)^{n+1/2}}{2} \\
&= \sum_{j \geq 0} [y^j] \left[(1 + \sqrt{1+y})^{j/2+n+1/2} \right] \frac{(1+y)^{n+1/2}}{2} \frac{1 - (-1)^j}{2} \\
&\quad - \sum_{j \geq 1} [y^{2j-1}] (1 - \sqrt{1+y})^{j+n} \frac{(1+y)^{n+1/2}}{2}.
\end{aligned}$$

Let us start with one term in the above sum:

$$\sum_{j \geq 0} [y^j] (1 + \sqrt{1+y})^{j/2+n+1/2} (1+y)^{n+1/2}.$$

This is of the form

$$\sum_{j \geq 0} [y^j] F(y) \Phi(y)^j$$

with

$$F(y) = (1 + \sqrt{1+y})^{n+1/2} (1+y)^{n+1/2} \quad \text{and} \quad \Phi(y) = \sqrt{1 + \sqrt{1+y}}.$$

This is the instance $x = 1$, which, by $\alpha = (1 + \sqrt{5})/2$, translates to

$$y = \frac{1 + \sqrt{5}}{2}, \quad F(\alpha) = \alpha^{4n+2}$$

and

$$\Phi'(\alpha) = \frac{3 - \sqrt{5}}{8}, \quad \frac{1}{1 - \Phi'(\alpha)} = 2 \left(1 - \frac{1}{\sqrt{5}} \right).$$

So our evaluation is:

$$2 \left(1 - \frac{1}{\sqrt{5}} \right) \alpha^{4n+2}.$$

The second term is

$$\sum_{j \geq 0} [y^j] (1 + \sqrt{1+y})^{j/2+n+1/2} (1+y)^{n+1/2} (-1)^j.$$

This is the instance $x = -1$, which translates to $y = -1$ and so the second term is

$$\frac{F(-1)}{1 + \Phi'(-1)} = 0.$$

Finally the last term is of the form:

$$\begin{aligned} & \sum_{j \geq 1} [y^{2j-1}] \left(1 - \sqrt{1+y}\right)^{j+n} (1+y)^{n+1/2} \\ &= \sum_{j \geq 1} [y^{2j-1}] y^{j+n} \left(\frac{1 - \sqrt{1+y}}{y}\right)^{j+n} (1+y)^{n+1/2} \\ &= \sum_{j \geq 0} [y^j] y^{n+1} \left(\frac{1 - \sqrt{1+y}}{y}\right)^{j+n} (1+y)^{n+1/2}. \end{aligned}$$

This is of the form:

$$\sum_{j \geq 0} [y^j] F(y) \Phi(y)^j$$

with

$$F(y) = \left(1 - \sqrt{1+y}\right)^n (1+y)^{n+\frac{1}{2}} y \quad \text{and} \quad \Phi(y) = \frac{1 - \sqrt{1+y}}{y}.$$

This is the instance $x = 1$, which translates to $y = \beta = \frac{1-\sqrt{5}}{2}$. Thus

$$F(\beta) = -\beta^{4n+2}, \quad \Phi'(\beta) = -\frac{1-\sqrt{5}}{4}, \quad \frac{F(\beta)}{1-\Phi'(\beta)} = -\left(1 + \frac{1}{\sqrt{5}}\right) \beta^{4n+2}.$$

So our evaluation is

$$\left[\left(1 - \frac{1}{\sqrt{5}}\right) \alpha^{4n+2} + \left(1 + \frac{1}{\sqrt{5}}\right) \beta^{4n+2} \right] \frac{1}{2} = F_{4n+1},$$

as claimed. □

Theorem 3. For $n > 0$,

$$\begin{aligned} F_{4n} &= \sum_{i=0}^n \sum_{j=0}^n \binom{n+i}{2j-1} \binom{n+j}{2i}, \\ F_{4n-3} &= \sum_{i=0}^n \sum_{j=0}^n \binom{n+i}{2j+1} \binom{n+j}{2i+1}. \end{aligned}$$

Again by using the Lagrange-Bürmann formula, Theorem 3 can be similarly proved.

Theorem 4. For $n > 0$,

$$\frac{F_{2n+2} + F_{n+1}}{2} = \sum_{0 \leq i, j \leq n} \binom{n-i}{2j} \binom{n-2j}{i}.$$

Proof. First, we replace i by $n-i$ and get

$$\sum_{0 \leq 2j \leq i \leq n} \binom{i}{2j} \binom{n-2j}{i-2j}.$$

Now we compute the generating function of it:

$$\begin{aligned} \sum_{n \geq 0} z^n \sum_{0 \leq 2j \leq i \leq n} \binom{i}{2j} \binom{n-2j}{i-2j} &= \sum_{0 \leq 2j \leq i} \binom{i}{2j} \frac{z^i}{(1-z)^{i+1-2j}} \\ &= \sum_{j \geq 0} \frac{z^{2j}(1-z)^{2j}}{(1-2z)^{1+2j}} = \frac{1-2z}{(1-z-z^2)(1-3z+z^2)} \\ &= \frac{1}{2} \frac{1}{1-z-z^2} + \frac{1}{2} \frac{1}{1-3z+z^2}, \end{aligned}$$

which is the generating function of the numbers $(F_{2n+2} + F_{n+1})/2$. \square

The following results are similar:

Theorem 5. For $n > 0$,

$$\begin{aligned} F_{2n} &= \sum_{i=1}^n \sum_{j=1}^n \binom{n-i}{j-1} \binom{n-j}{i-1}, \\ F_{2n-1} &= \sum_{0 \leq j \leq i \leq n} \binom{n}{i-j} \binom{n-i}{j}. \end{aligned}$$

Theorem 6. For $n > 0$,

$$F_{2n} + 1 = \sum_{i=0}^n F_{2i-1} = \sum_{0 \leq i \leq j \leq n} \binom{n-i}{j} \binom{j}{2i}. \quad (2.1)$$

Proof. Multiplying the right hand side of (2.1) by z^n and summing over n , we get

$$\begin{aligned} S &= \sum_{n \geq 0} z^n \sum_{0 \leq i \leq j \leq n} \binom{n-i}{j} \binom{j}{2i} = \sum_{0 \leq i \leq j} \sum_{h \geq 0} z^{h+i+j} \binom{h+j}{j} \binom{j}{2i} \\ &= \sum_{0 \leq i \leq j} \binom{j}{2i} z^{i+j} \sum_{h \geq 0} z^h \binom{h+j}{j} = \sum_{0 \leq 2i \leq j} \binom{j}{2i} z^{i+j} \frac{1}{(1-z)^{j+1}} \\ &= \sum_{i \geq 0} \frac{z^{3i}}{(1-2z)^{2i+1}} = \frac{1-2z}{(1-z)(1-3z+z^2)} = \frac{z}{1-3z+z^2} + \frac{1}{1-z}, \end{aligned}$$

which is the generating function of the numbers $F_{2n} + 1$. \square

For the Pell numbers, we give the following result:

Theorem 7. For $n \geq 0$,

$$P_{n+1} = \sum_{0 \leq i \leq j \leq n} \binom{n-i}{j} \binom{j}{i}. \quad (2.2)$$

Proof. Multiplying the right hand side of (2.2) by z^n and summing over n , we get

$$\begin{aligned} S &= \sum_{n \geq 0} z^n \sum_{0 \leq i \leq j \leq n} \binom{n-i}{j} \binom{j}{i} = \sum_{0 \leq i \leq j} \sum_{h \geq 0} z^{h+i+j} \binom{h+j}{j} \binom{j}{i} \\ &= \sum_{0 \leq i \leq j} \binom{j}{i} z^{i+j} \sum_{h \geq 0} z^h \binom{h+j}{j} = \sum_{0 \leq i \leq j} \binom{j}{i} z^{i+j} \frac{1}{(1-z)^{j+1}} \end{aligned}$$

$$\begin{aligned}
&= \sum_{0 \leq i \leq j} \frac{z^j}{(1-z)^{j+1}} \binom{j}{i} z^i = \sum_{j \geq 0} \frac{z^j}{(1-z)^{j+1}} (1+z)^j \\
&= \frac{1}{1-z} \frac{1}{1 - \frac{z(1+z)}{1-z}} = \frac{1}{1-2z-z^2}.
\end{aligned}$$

This is the generating function of the numbers P_{n+1} . □

Now we give a double sum for the Tribonacci numbers:

Theorem 8. For $n \geq 0$,

$$T_n = \sum_{0 \leq j \leq i \leq n} \binom{n-i}{i-j} \binom{i-j}{j}.$$

Proof. Consider

$$\begin{aligned}
\sum_{n \geq 0} T_n z^n &= \sum_{0 \leq j \leq i \leq n} z^n \binom{n-i}{i-j} \binom{i-j}{j} = \sum_{0 \leq j \leq i} z^i \binom{i-j}{j} \sum_{h \geq 0} z^h \binom{h}{i-j} \\
&= \sum_{0 \leq j \leq i} z^i \binom{i-j}{j} \frac{z^{i-j}}{(1-z)^{i-j+1}} = \sum_{j \geq 0} \sum_{h \geq 0} z^{h+j} \binom{h}{j} \frac{z^h}{(1-z)^{h+1}}.
\end{aligned}$$

Let $t = \frac{z^2}{1-z}$, and we continue

$$\begin{aligned}
\sum_{n \geq 0} T_n z^n &= \frac{1}{1-z} \sum_{0 \leq j} z^j \sum_{h \geq 0} \binom{h}{j} t^h = \frac{1}{1-z} \sum_{0 \leq j} z^j \frac{t^j}{(1-t)^{j+1}} \\
&= \frac{1}{1-z} \frac{1}{1-t} \frac{1}{1 - \frac{zt}{1-t}} = \frac{1}{1-z} \frac{1}{1-t-zt} \\
&= \frac{1}{1-z} \frac{1}{1 - \frac{z^2}{1-z} - \frac{z^3}{1-z}} = \frac{1}{1-z-z^2-z^3},
\end{aligned}$$

which is the generating function of the Tribonacci numbers, as expected. So the proof is complete. □

By using the same proof method as in Theorem 8, we get a more general result:

Theorem 9. For $n > 0$,

$$f_n^{(k)} = \sum_{0 \leq i_k \leq \dots \leq i_1 \leq n} \binom{n-i_1}{i_1-i_2} \binom{i_1-i_2}{i_2-i_3} \dots \binom{i_{k-1}-i_k}{i_k}$$

where $f_n^{(k)}$ is the n -th generalized order- k Fibonacci number.

REFERENCES

- [1] G. E. Andrews, R. Askey and R. Roy, Special functions. Encyclopedia of Mathematics and its Applications, 71. Cambridge University Press, Cambridge, 1999.
- [2] A. T. Benjamin and J. J. Quinn, Proofs That Really Count, Mathematical Association of America, 2003.
- [3] J. Ercolano, Matrix generator of Pell sequence, Fibonacci Quart. 17 (1) (1979), 71–77.
- [4] P. Henrici, Applied and computational complex analysis. Vol. 1. John Wiley & Sons, Inc., New York, 1988.

- [5] A. F. Horadam, Pell identities, *Fibonacci Quart.* 9 (3) (1971), 245–252, 263.
- [6] R. Knott, Fibonacci and Golden Ratio Formulae, <http://www.maths.surrey.ac.uk/hosted-sites/R.Knott/Fibonacci/>.
- [7] J. Riordan, *Combinatorial Identities*, John Wiley & Sons, Inc., New York, 1968.

TOBB UNIVERSITY OF ECONOMICS AND TECHNOLOGY, MATHEMATICS DEPARTMENT, 06560
ANKARA TURKEY

E-mail address: ekilic@etu.edu.tr

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH 7602 STELLENBOSCH SOUTH
AFRICA

E-mail address: hproding@sun.ac.za